


# 2013 WINTER WHEAT VARIETIES

## Performance Evaluation and Recommendations

Recommendations are made for the districts shown on the map below



by the Montana State University  
Agricultural Experiment Station

The information in this publication can also be found at a link on:

<http://plantsciences.montana.edu/crops>

Another variety selection tool is available at :

<http://www.sarc.montana.edu/php/varieties.html>

## 2013 Recommended Varieties: Hard Winter Wheat and Soft White Winter Wheat for Montana by District

Variety	Districts (see map on cover)					
	1 Northwest	2 Southwest	3 Southeast	4 Central	5 North Central	6 Northeast
<b>Hard Red and Hard White Winter Wheat</b>						
Bearpaw ++ <sup>2/</sup>			D	D	D	
Bynum (P) <sup>2/</sup> +				D	D	
Carter (P)+		D	D	D	D	D
CDC Falcon (P)+		DI	DI	DI	DI	DI
Colter (MT08172)++		D	D	D	D	
Decade +			D	D	D	D
Genou + <sup>2/</sup>			D	D	D	
Hyalite (HWW, P)+		D	D	D	D	
Jagalene (P)+	D	D	D	D	D	
Jerry						D
Judee ++ <sup>2/</sup>			D	D	D	
Ledger (P)+		D		D	D	
Norris (P)+		D	D	D		
Promontory <sup>1/</sup>	D	D	DI	D		
Pryor (P)+		D	D	D	D	D
Rampart <sup>2/</sup>			D	D	D	
WB-Quake (P)++	D	D	D	D	D	D
Warhorse (MTS0808)++ <sup>2/</sup>			D	D	D	
Yellowstone +	D	D	D	D	D	
<b>Soft White Winter Wheat</b>						
Eltan	D	D				
Hill 81	D	D				
Lewjain	D					
Malcolm	D	D				

HWW = Hard White Winter Wheat

D = Dryland

I = Irrigated

(P) = a Private Variety

+ = a "Protected" variety under the Plant Variety Protection Act

++ = PVP Title V pending

<sup>1/</sup> = dwarf smut resistant

<sup>2/</sup> = sawfly areas only

## TABLE OF CONTENTS

	<u>Page</u>
Hard Red Winter and Soft White Winter Wheat Varieties Recommended by the Montana Agricultural Experiment Station .....	Inside Cover
Introduction.....	1
Variety Testing Procedures .....	1
Description of Data Collected .....	1
Table 1. Summary of Agronomic Practices .....	2
Statistical Analyses and Interpretation .....	3
2012 Test Conditions.....	3
Dwarf Smut (TCK) .....	4
What Recommendation by MAES Means .....	4
Producing Winter Wheat.....	5
Yield in Winter Wheat as Influenced by Percent Stand .....	6
Hard Red Winter Wheat Comparisons:	
Table 2. List of Varieties .....	7
Table 3. District 1 - Kalispell - Dryland (High Rainfall) .....	8
Table 4. District 2 - Bozeman - Dryland .....	9
Table 5. District 3 - Huntley - Dryland .....	10
Table 6. District 4 - Moccasin - Dryland .....	11
Table 7. District 5 - Conrad - Dryland.....	12
Table 8. District 5 - Havre - Dryland.....	13
Table 9. District 6 - Sidney - Dryland.....	14
Table 10. Williston, North Dakota - Dryland .....	15
Table 11. Yield in winter-kill environments.....	16
Table 12. Yield performance under sawfly pressure .....	17
Table 13. Precipitation and average monthly temperature for Crop Year .....	18
Table 14. Selected agronomic characters, cereal quality evaluations and disease reactions .....	19
Additional Descriptive Information for Winter Wheat Varieties:	
Hard Winter Wheat .....	20
Plant Variety Protection .....	23
Acknowledgements .....	24


# WINTER WHEAT VARIETY PERFORMANCE SUMMARY IN MONTANA

J. E. Berg, P. L. Bruckner, G.W. Bergman, B. Deanon, J. Eckhoff, K. D. Kephart, P. Lamb, S. Loomer  
J. H. Miller, M. Peterson-Walter, G.V.P. Reddy, R.N. Stougaard, D.M. Wichman, A. Dyer, W. Grey, D. Nash,  
and R. Larson

## Introduction

The agronomic characteristics of winter wheat varieties recently developed or evaluated by the Montana Agricultural Experiment Station are compared in this publication with other varieties grown in the state. Varieties recommended for production in the respective districts of Montana are designated by an **R**. A brief description of each variety is given which may include a variety's particular advantages or disadvantages. The information was extracted from the Intrastate Winter Wheat Nursery. This data is prepared by research personnel of the Montana Agricultural Experiment Station. Where available, up to four years of yield data are shown for the varieties. In some years data are not available because of hail, frost, or other unavoidable causes.

## Variety Testing Procedures



**Fig. 1. Test Locations for Montana winter wheat performance tests in 2012.**

## Entries

Names of commercially available entries evaluated in 2012 are listed with their origins, experimental designation, release year, and pedigrees in Table 2 for the hard winter wheats and in Table 15 for the soft white wheats. Forty-nine hard wheats are included in this summary comprising 30 varieties (17 public and 13 private) and 19 experimental lines (all public). Numbered entries preceded by a state designation [e.g. MT0978 (Montana), MTS1024 (Montana)] are experimental lines provided by the breeder.

## Experimental Design and Seeding Methods

The Intrastate Winter Wheat Test consisted of a 49 entry test with 3 replicates. It was planted in the form of 7x7 lattice at all. Plot size varied by location, from 35 ft<sup>2</sup> at Conrad to 60 ft<sup>2</sup> at Havre. Row number varies: Bozeman and Havre are 3-row, Conrad, Huntley, and Sidney are 4-row, Moccasin (5-row), Kalispell (7-row), and Williston (8-row) Row spacing at all locations was on 1 ft. centers, except at Williston and Kalispell (6" centers). All plots were seeded at 0.6 grams seeds/ft<sup>2</sup>, which is roughly equivalent to 1 bushel per acre, except at Williston where the seeding rate was about 77 pounds per acre. Information on previous crop, planting date, fertilizer use and harvest date is available in Table 1.

All seed for each nursery was treated with Cruiser Maxx Cereals seed treatment at recommended rates before planting.

## Description of Data Collected

### Yield

All rows of each plot were trimmed and measured and harvested using an experimental plot combine. Grain yields are reported in bushels per acre based on a 60 pound standard bushel weight. In addition to yields obtained in 2012, data is provided for two (2011-2012), three (2010-2012) and four (2009-2012) year averages for hard wheat entries tested during previous cropping seasons

### Locations

Hard winter wheats were planted at 7 Montana and 1 North Dakota location (Fig. 1) including Conrad and Havre in the North Central district, Moccasin in the Central district, Huntley in the Southern district, Sidney and Williston, ND representing the Northeast district, Kalispell in the Northwest and Bozeman in the Southwest districts of the state.

**Table 1. Summary of agronomic practices used on hard winter wheat performance trials in Montana in 2012. Fall nitrogen (N), phosphorus (P<sub>2</sub>O<sub>5</sub>) and potassium (K<sub>2</sub>O) were preplant applied and incorporated.**

Location	2011 Crop	2010 Crop	2011 Planting Date	Fertilizer				2012 Harvest Date
				N		P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	
				Fall	Spring			
----- Pounds per acre -----								
Kalispell	peas	barley	Sep 24	10	100	35	90	Aug 14
Bozeman	fallow	spring wheat	Oct 2	157	-	0	0	Aug 3
Huntley	chem. fallow	fallow	Sep 26	60	-	30	0	Jul 18
Moccasin	chem. fallow	barley	Sep 27	10	60	10	10	Jul 26
Conrad	fallow	barley	Sep 22	41	115	0	0	Aug 22
Havre	fallow	spring wheat	Sep 30	70	-	40	25	Aug 2
Sidney	fallow	safflower	Oct 13	40	-	0	0	Jul 31
Williston, ND	fallow	safflower	Sep 23	75	-	26	0	Jul 25

### Test Weight

Test weight (pounds per bushel) were obtained for each plot by using Dickey-John Grain Analysis Computer (GAC) at some locations. Other locations use a Seedburo test weight apparatus. In this case, a sample is dropped through a funnel at a given height into a quart brass bucket, excess grain is removed by a flat stick then weighed on a gram scale, and grams per quart are converted into pounds per bushels.

### Heading Date

Heading date is taken when 50% of the heads in a plot were extended above the flag leaf collar. Heading dates are recorded both in ordinal date (number of days from January 1) and the actual calendar date.

### Plant Height

Plant height was measured in inches from the soil surface to the top of the head, excluding the awns.

### Grain Protein

Grain protein is sampled from a composite of all 3 replicated plots at each location. It is determined as a % by NIR (near infrared reflectance) on the Infratec whole grain analyzer. Samples are adjusted to a 12% moisture basis.

### Winter Survival

Percent winter survival is estimated for each plot after initial spring green-up at locations where

significant winter injury occurred. In 2012, Williston had 22% winter survival noted on June 6<sup>th</sup> [range = 5 (Judee) – 57% (Peregrine)].

Table 11 contains information on % winter survival and associated yield in winter-kill environments from 2005 to 2012. The data summarizes 10 tests in which significant winter-kill occurred (test average for winter survival was less than 90%). All sites with winter-kill were in District 6 (Sidney and Williston) which is the most severe location for winter wheat survival of our testing locations.

### Wheat Stem Sawfly

Wheat stem sawfly (WSS) is a persistent and economic problem for wheat growers in Montana. Currently, Montana wheat acreage infested by WSS is primarily in the north central (District 5), central (District 4) and south central (District 3) cropping districts. Host plant resistance in the form of stem solidness has been effective in reducing sawfly losses in both spring and winter wheat. Solid-stemmed winter wheats, 'Vanguard' (dropped from testing in 2009) and 'Rampart' were released in 1995 and 1996, respectively. These 2 varieties were planted on 6% of the winter wheat acreage in the 2012 crop year (Rampart was the leading variety planted in the 2003 to 2006 crop years). Both these varieties have marginal winter hardiness. 'Genou', released in 2004, was the leading variety during the period 2007-2011. In 2012, Genou was planted on 20% of the winter wheat acreage (second in acreage to Yellowstone at 23%). Newer solid-stemmed varieties include Bearpaw (2011), Judee (2011), and WB-Quake (2010).

## Statistical Analyses and Interpretation

Table 12 contains information on yield and % sawfly cutting at 17 testing locations where sawfly pressure was present during the years 2005-2012. The data is from Havre, North Havre (a site 25 miles north of Havre), Loma (15 miles northeast of Ft. Benton), Turner (60 miles east-northeast of Havre), and Willow Creek (35 miles west-northwest of Bozeman). Solidness scores (rated on a 5-25 scale) are shown for solid and semi-solid varieties in Table 14.

### Coleoptile Length

Coleoptile length evaluation was performed in Bozeman under controlled (growth chamber) conditions. Twenty-five seeds per variety were planted in wetted vermiculite. After 15 days the coleoptile (sheath covering the emerging shoot that helps penetration to the soil surface) was measured. This test was replicated 3 times for each variety. Results from previous years are reported in Table 14. Long coleoptiles are generally longer than 4 inches, medium from 2.7-4 in, and short are under 2.7 in. Care should be taken not to plant short coleoptile varieties too deep.

### Other Agronomic Characters

Table 14 contains information on grain maturity, chaff color, relative winter survival and straw strength for the hard wheat varieties listed in this publication.

### Cereal Quality

Milling and baking characteristics for varieties are presented in Table 14. They are rated for each variety on a 1-5 scale (5 = superior). A quantitative polyphenol oxidase (PPO) has been determined for varieties since the 2006 mill and bake evaluation. These varieties are reported in Table 14 as low to high. A lower value is associated with better Asian noodle quality.

### Disease Reactions

Disease reactions for hard red wheat varieties are listed in Table 14. There is information on dwarf smut, stripe rust, stem rust and leaf rust. Table 18, for soft white winter wheat, contains information on dwarf smut, snow mold, stem rust and stripe rust.

The data collected at each winter wheat location was analyzed as a three-replication lattice or randomized complete block design. Least significant difference at the 0.05 probability level (LSD,  $p = 0.05$ ) and coefficients of variation (CV) were calculated from analysis of variance at each location. The LSD is used to compare the performance of two specific varieties at a time. If the difference between two varieties exceeds the LSD this is interpreted as a true difference, because a difference between two varieties this large will only occur 5% of the time due to chance.

Tables 3 through 10 show 2012 data for hard winter wheat collected at all harvested experiment station sites. Where a variety has been in the test for two, three or four years, combined analyses of the yield data over years are presented.

Variety selection should be based on yield stability at a particular location over a period of years. Selection should also consider test weight, winter-hardiness, heading date, plant height, protein and disease resistance.

## 2012 Test Conditions

Statewide winter wheat yields were projected by the Montana Agricultural Statistics Service at 38 bu/a for 2012 compared to 41 bu/a for the 2011 harvest year. The harvested acreage in 2012 was 2.14 million acres (total production = 81.3 million bu) compared 2.19 million acres in 2011 (total production = 89.7 million bu). Rainfall for the 2011-2012 winter crop year was below average at all locations tested (Table 13), except Kalispell. Average yearly temperatures were above long term at all locations, ranging from +0.1° at Conrad to +7.3°F at Huntley.

Test weight averaged 58.8 lb/bu across all locations. Conrad (61.2) and Huntley (63.2) were the only locations above 60 lb/bu.

Winterkill at Williston, 22% survival across varieties, reduced yields of susceptible varieties (Art, AP503 CL2, Bynum, Curlew, Genou, Judee, Ledger, Promontory, and SY Wolf). No winterkill was recorded at Sidney.

Heading dates were earlier in 2012 for Bozeman (2 days), Sidney (7 days), Williston (5 days), and Huntley (4 days) than long term averages. Havre

(2 days) and Kalispell (6 days) had later than average heading dates. Conrad was average and Moccasin had no data.

Stripe rust (average = 79%, range 36 – 100% on July 12) at Kalispell was a significant factor in yield (range 9 – 108 bushels per acre) reduction for highly susceptible varieties (Accipiter, Bearpaw, Broadview, Carter, CDC Falcon, Decade, Genou, Jerry, McGill, Norris, overland, and Pryor). There was some sawfly cutting recorded at the Havre Experiment Station averaging 8% of stems cut across varieties (range = 0 – 20%).


Protein content averaged 13.1% across all locations (location range = 11.0 – 15.2%) tested. Sidney, Moccasin, Huntley, and Conrad were below 12%. The range of genotype means across all locations was 12.3 (Pryor) – 14.2 (Genou) %.

Leading winter wheat varieties planted for 2012 were Yellowstone (22.8%), Genou (19.7%), CDC Falcon (7.0%), Jerry (4.9%), Ledger (4.9%), and AP503 CL2 (4.5%).

### Dwarf Smut (TCK)

Dwarf smut (TCK) can be controlled with 'Dividend' seed treatment (see page 5). Dwarf smut or dwarf bunt (*Tilletia controversa* Kuhn) is a fungal disease that occurs in areas where winter wheat is subjected to prolonged snow cover or unfrozen ground. The planting of dwarf smut resistant varieties (Promontory and Lewjain are currently recommended) as one practical means of control.

The amount of wheat lost each year because of dwarf smut is small in relation to the state's total crop, but individual operators may experience severe losses in heavily infested, localized areas.



**Fig. 2. Known areas of dwarf smut (TCK) infestations.**

If you farm in the vicinity of one of the shaded areas in the map (Figure 2.), you would be well advised to observe closely your winter wheat crop and consider planting a resistant variety (Tables 14 and 18) or use 'Dividend' seed treatment, only.

### What Recommendation by MAES Means

Classification of winter wheat varieties is determined on a yearly basis by the Montana Agricultural Experiment Station (MAES) Wheat Variety Release Committee. This 16 member committee is composed of one wheat breeder, one cereal or forage quality scientist, one plant pathologist, one entomologist, one weed scientist, one cropping systems specialist, six Research Center agronomists, one manager from both the Montana Foundation Seed program and the Montana Seed Growers Association, one Montana Wheat and Barley Committee member and one representative of the Montana Agricultural Experiment Station Advisory Board.

A variety is eligible for recommendation when a minimum of 16 location-years of performance data is obtained from the Montana State University statewide winter wheat performance trials. Test results indicate that the variety is equal to or superior in overall merit to specified check cultivars and has end-use quality equal to or exceeding currently recommended varieties. For varieties originating from private companies, recommendation is considered only at the request of the company when adequate data is available.

Recommendations of varieties are considered on a case by case basis. Yield performance of a variety is an important criteria, but also considered are test weight, grain protein content, winter survival, pest resistance and end-use quality data. In general, yield needs to be at least equal to currently recommended varieties in a particular district, unless the variety is being recommended for a specific purpose, e.g. winter hardiness, sawfly resistance. For example, Rampart, which is not competitive in the absence of wheat stem sawfly, is recommended in Districts 3, 4 and 5 for sawfly areas only. Only six varieties are recommended for the Northeast district due to severe winter conditions and a higher probability of stem rust in this region. Thus varieties recommended for District 6 must have higher winter survival and stem rust resistance.

If a serious defect in the variety is identified during performance testing, the variety will not be recommended. Examples of defects resulting in non-recommendation include: high probability of winter-kill, low grain protein, low baking quality, etc.

Lack of variety recommendation by MAES may occur due to a decision by the originating company not to test the variety in statewide performance trials. In this case the lack of recommendation is due to inadequate or no data rather than a specific varietal defect.

Montana produces primarily hard red winter and hard red spring wheats. Continuous improvement of the milling and/or baking quality of Montana grown winter wheat is one of many objectives of the Montana Agricultural Experiment Station breeding and cultivar development program. All varieties recommended by the Montana Agricultural Experiment Station have been evaluated and found to be acceptable for milling and baking performance by the Cereal Quality Laboratory at Montana State University.

The quality of Montana recommended varieties, if grown and marketed within their respective classes, is acceptable by domestic users. Montana's future as a hard red and hard white winter wheat producing state for both the domestic and export markets rests on the quality of the product.

## Producing Winter Wheat

Plant **CERTIFIED CLASS SEED** of varieties **RECOMMENDED** by the Montana Agricultural Experiment Station.

### Seed Treatment

Treat all winter wheat seed with a recommended fungicide to reduce losses caused by cereal smut or other seed-borne diseases. Several non-mercurial compounds are registered for grain seed treatment.

Dwarf smut (bunt) can be controlled with difenoconazole. Dividend® contains this compound and is available in Montana. If you farm in a dwarf smut area contact your seed dealer or chemical representative for more information about this seed treatment. See page 4 for known areas of dwarf smut infestations.

Diseases are best controlled when all seeds are coated with a seed treatment. Do not over-treat-- Follow recommendation of manufacturer of product as to rate.

Truck-mounted seed treaters, which apply the fungicide as the seed is augered into the drill box, do a good job of treating if operated according to manufacturer's specifications.

Drill box treatments are not effective for general use.

When using any pesticide materials, read the information on the label as to rate of application, specific uses, methods of handling, precautions, etc.

### Seeding Rate and Date

The following rates and dates for seeding are general (Figure 3). The heavier seeding rate, where indicated, is applicable to plump seed of high test weight (above 60 lbs/bu) or for seed having a kernel size larger than normal for most other varieties. The lighter rates are for the smaller seeded varieties or when test weight is below normal for larger seeded varieties. Seeding rates may be lower if adequate nitrogen and phosphorus amounts are applied at planting.

Winter wheat seed lots may vary in the number of seeds per pound depending on the ratio of large-to-small seeds in a seed lot. The average is approximately 15,000 seeds per pound. A precise count of the number of seeds per pound should be made on your seed lot to help calibrate your drill. You can also calculate how many pounds of seed you will need to plant an acre.

**Figure 3. Seeding rate and date for winter wheat**

Districts	Dryland	Irrigated	Date of Seeding
5,6	30-60	60-75	Sept. 1-15
1,2,3,4	30-60 (10-20 seeds/sq. ft.)	60-75 (20-25 seeds/sq. ft.)	Sept. 10-25

As to seeding date -- DO NOT SEED TOO EARLY in areas where root rot diseases are prevalent. In areas where Cephalosporium stripe, wheat streak



mosaic virus or other root rot diseases have caused losses, delay seeding until the soil temperature in the seed zone will stay below 55°F except for brief periods during the day. In the southern half of Montana, this is usually September 10 to 20. In Districts 5 and 6, seed between September 1 and 15. Cooler soil temperatures slow root development and reduce the probability of winter root injury and invasion by soil-borne organisms. To reduce the incidence of root and foot rots, plant winter wheat on land previously seeded to other crops such as barley, oats or spring wheat. Extreme seeding delay, however, reduces seedling vigor and increases chances of winter-kill.

### **Seeding Depth**

Set the drill to place the seed 1 to 2 inches below the soil surface. Deeper seeding reduces tillering and lowers crop yields. With the furrow drills, wind-driven soil particles settle in the furrows covering the seed deeper than desired.

<p><b>Yield in Winter Wheat as Influenced by Percent Stand</b></p>
--

During periods of winter injury farmers are frequently faced with a decision as to whether or not a field should be torn up and re-seeded. A 40 to 50 percent winter wheat stand, if general over field, may produce as much as re-seeded spring wheat. Thinner stands will likely demand more attention for weed control.

The guidelines for evaluating winter wheat stands are to determine the average number of healthy plants per square yard. We suggest making a square frame out of 3/8 inch rod. Walk the field in a zigzag pattern counting at ten random locations.

Fields that have 80 or more plants per square yard will probably produce more than if replanted to spring wheat (information taken from 1995 Master's Thesis, "Critical Overwintering Plant Population for Successful Winter Wheat Production in Montana" by Doug Holen).

**Table 2. List of public and private hard winter wheat varieties.**

Variety	Experimental Designation	Origin	Release Year	Pedigree
<b>Public Varieties</b>				
Accipiter	DH00-18-196	Saskatchewan	2008	CDC Raptor/CDC Falcon
Bearpaw	MTS0721	Montana	2011	selection from a composite of 5 crosses: 99X96, DMS/Rampart//Pronghorn/3/2*/Rampart; 99X97, DMS/Rampart//Pronghorn/3/ Rampart/4/ (MTW9806, Redwin/Rio Blanco/NuWest) ; 99X98, DMS/Rampart//Pronghorn/3/Rampart/4/NuPlains; 99X99, DMS/Rampart//Pronghorn/3 /Rampart/4/(MT9513, NuWest/5/(TAM W-103/Froid/4/Yogo/Turkey Red/3/Centurk, MT8030)); and 99X100, DMS/Rampart//Pronghorn/3/ Rampart/6/(MT98113, Judith/5/ (MT8764, Crest/(VT1230, French male sterile line)/4/((PI178383/ Cheyenne/3*Tendoy, ID5011)/3/(ID5006, Norin 10/Staring// 2*Cheyenne), ID745101)))
Broadview	LE1911	Alberta	2009	KS92WGRC15/CDC Kestrel//CDC Falcon
Colter	MT08172	Montana	2013	(Yellowstone sib, MT9982)*2/(BZ9W96-895, ped. unknown from male sterile pop.)
Curlew	UT9325-55	Utah	2009	Golden Spike sib/3/Manning/R-82-1859//Weston
Decade	MT0552	Montana; North Dakota	2010	selection from composite of 3 crosses:(Sumner sib, KS831936-3, (Plainsman V/Odesskaya 51)/(NE86501, Colt/Cody), N95L159, Wesley sib)/3/ CDC Clair, N95L159/(MT9602, NuWest/Tiber) and N95L159/4/ (MT9609, Froid/SD1287// Redwin/3/NuWest)
Genou	MTS0031	Montana	2004	(Lew/Tiber//Redwin, MTS92015)/3/Vanguard/ Norstar
Jerry	ND9257	North Dakota	2001	Roughrider/(ND7571, Winoka/NB66425)/3/ Arapahoe
Judee	MTS0713	Montana	2011	(Vanguard/Norstar/Judith dwf, 93X312E14)/3/ NuHorizon
McGill	NE01481	Nebraska	2010	((Vona//Chisholm/Plainsman V, OK83201)/3/ Redland, NE92458)/4/Ike
Overland	NE01643	Nebraska, South Dakota	2007	(Millenium sib, NE94482)/(ND8974, Seward/ Archer)
Peregrine	DH99-37-100	Saskatchewan	2008	McClintock/S86-808
Promontory	UT1567-51	Utah	1990	Manning/Bezostaya-1
Rampart	MTS92042	Montana	1996	Lew/Tiber//Redwin
Robidoux	NI04421	Nebraska	2010	(Odesskaya polukarlikovaya/Cody//Pavon/ 3*Scout 66, NE96644)/3/Wahoo sib
Warhorse	MTS0808	Montana	2013	selection from a composite of 3 crosses: 00X182, ((Froid/Winoka/7/ ((Sinvalocho/Wichita// Hope/Cheyenne /3/Wichita/4/Seu Seun 27, TX55-391-56-D8)/5/Westmont, MT6928)/6/ Trader, MT85200)/8/ Redwin, MT9908)/9/ NuPlains/6/(MTS9862, (NuWest/ Lovrin 24 /4/((Rego/ Cheyenne, Sel. 39-18-7)/ Winalta, MT7431)/3/(MT7115, Yogo/T. polonicum-70-5), MT91366)/5/ (MTS92137, Lew/Tiber//Redwin)); 00X183, NuPlains/MTS9862/4/ (MTW0047, Judith/(PI262605, Karagach, RWA resis.)/3/(S86-740, Norstar/ Plainsman V //Ulianovka)); and 00X184, NuPlains/MTS9862/5/(MTS0028, Vanguard/4/(Lew/Tiber// Redwin, MTSF1570)/3/ Norstar)
Yellowstone	MT00159	Montana	2005	F2 composite of Promontory/Judith and Judith-dwarf/Promontory
<b>Private Varieties</b>				
AP503 CL2	CL03040-5-2	AgriPro	2007	iW98-362A1 (Als3-653)/AP502 CL (Als1-653) [CLEARFIELD] (Note: W98-362 = Jagalene)
Art	98x0338-13	AgriPro, Sygenta	2007	Jagger/4/(W94-244-132, (TAM 200/ Mesa sib, W189-088)/3/(W188-052, (C78-244/Archer, 82F2042#2)// Mesa sib))
Bynum (CL)	MTCL0318	WestBred LLC, Montana	2005	Rampart/FS2//CDC Kestrel, FS2 = mutagenized Fidel
Carter	BZ9W02-2060	WestBred LLC	2006	Jagger/Rampart
CDC Falcon	S94-4	Western Plant Breeders/Saskatchewan	1999	Norstar*2/Vona//Abilene
Jagalene	W98-362	AgriPro Seeds	2002	Jagger/Abilene
Ledger	BZ9W96-788-d	WestBred LLC	2004	(Hatten/SS-14, BZ9W92-709)/3/(MTSF1142, Lew/ Tiber//Redwin)
Norris (CL)	MTCL0316 (IMI)	WestBred LLC, Montana	2005	Big Sky/(TXGH 12588-26, TAM-110 sib)*4/FS2
Pryor	BZ9W96-919	WestBred LLC	2002	Hatten/Abilene
Radiant	W337	Alberta, Meridian Seeds	2002	Norstar*6/Cmc1//Norwin/UT125512, WSMV resistant
SY Clearstone 2CL	MTCL1077	Sygenta, Montana	2012	Yellowstone*4/3/MTCL01158/CDC Teal 11A//Jagalene
SY Wolf	BC01007-7	AgriPro, Sygenta	2010	W99-331/97x0906-8
WB-Quake	BZ9W05-2043	WestBred LLC (Monsanto)	2011	Rampart/Kestrel

Table 3. HARD WINTER : District 1-- Kalispell - Dryland (High Rainfall)

Cultivar/Line	Grain Yield (bushels/acre)				2012 Data					
					Test weight	Heading Date		Plant height	Stripe rust	Protein
	2012	2011-12	2010-12	2009-12		Ordinal	Calendar			
		2 yr	3 yr	4 yr	lb/bu	from Jan1		in	%	%
Accipiter ++	21.9	36.9	70.6	75.8	53.0	171.7	20-Jun	40.4	97.5	12.6
AP 503 CL2 (P, CL)+	47.6	73.3	97.0	93.0	55.2	166.6	15-Jun	40.4	70.0	13.8
Art (P)+	44.9	53.4	84.2		47.4	161.7	10-Jun	41.0	97.5	14.4
Bearpaw	14.9	25.8	67.6	70.0	47.8	168.7	17-Jun	40.6	100.0	16.8
Broadview	12.0	18.7	56.7		50.3	171.3	19-Jun	38.9	99.7	14.0
Bynum (P, CL)+	48.8	63.5	88.0	83.7	<b>58.4*</b>	164.3	12-Jun	45.9	79.7	14.0
Carter (P)+	10.3	14.6	54.9	61.9	48.2	169.0	17-Jun	35.2	99.5	16.1
CDC Falcon (P)+	38.8	45.6	76.9	80.4	51.2	166.7	15-Jun	38.1	98.4	13.1
Curlew <sup>1/</sup>	73.3	93.2	<b>114.9*</b>	<b>111.5*</b>	<b>58.0*</b>	165.0	13-Jun	43.7	91.5	13.9
Decade +	9.1	12.8	51.1	60.3	44.3	165.3	13-Jun	40.1	99.1	15.3
Genou +	16.1	21.1	60.6	65.3	50.4	171.4	19-Jun	42.6	100.0	15.9
R Jagalene (P)+	64.5	66.6	95.5	<b>95.8*</b>	55.9	165.6	14-Jun	41.8	84.6	14.1
Jerry	15.5	21.9	53.5	61.5	49.1	171.4	19-Jun	44.6	98.3	13.8
Judee ++	60.7	83.0	104.6	<b>99.5*</b>	52.4	167.3	15-Jun	40.9	<b>53.7*</b>	14.6
Ledger (P)+	43.6	45.5	76.9	79.7	56.5	165.3	13-Jun	40.6	97.4	13.0
McGill +	20.8	38.1			46.8	166.0	14-Jun	40.6	98.0	13.8
MT08172 (Colter)++	<b>95.3*</b>	<b>108.6*</b>			<b>59.1*</b>	170.3	18-Jun	44.0	61.6	13.2
MT0871	52.9	72.6	102.5		51.8	171.7	20-Jun	43.1	72.9	15.3
MT0978	60.2	81.0			53.9	169.6	18-Jun	41.5	73.6	14.8
MT10113	12.3				48.0	164.4	12-Jun	39.2	99.9	16.3
MT10116	84.1				<b>57.4*</b>	170.7	19-Jun	42.1	60.1	12.9
MT1078	63.0				55.8	167.0	15-Jun	42.0	72.4	14.5
MT1088	77.0				57.1	169.0	17-Jun	41.2	64.4	13.1
MT1090	83.2				56.7	168.0	16-Jun	44.4	75.4	12.7
MT1091	72.4				55.0	168.7	17-Jun	42.7	75.1	13.4
MT1092	84.4				<b>57.3*</b>	169.7	18-Jun	44.1	56.1	12.4
MT1105	64.8				56.1	168.7	17-Jun	41.9	75.8	13.1
MT1155	50.3				53.2	169.0	17-Jun	41.1	83.5	14.5
MT1156	73.4				55.5	169.7	18-Jun	42.8	57.9	13.2
MTCL1067	55.6	73.9			54.4	167.0	15-Jun	46.4	68.3	14.1
MTS0808 (Warhorse)++	74.6	94.4	<b>113.9*</b>		<b>58.5*</b>	168.4	16-Jun	38.8	<b>51.4*</b>	14.3
MTS0819	36.2	57.0	89.6		47.4	169.0	17-Jun	39.4	88.3	14.5
MTS0819-98 (HWW)	74.0				53.6	168.7	16-Jun	38.6	<b>48.1*</b>	14.5
MTS1024	67.6				56.6	169.0	17-Jun	39.8	62.0	14.3
MTW08168	97.6*	<b>107.9*</b>			<b>58.8*</b>	172.0	20-Jun	47.5	<b>49.0*</b>	12.9
Norris (P, CL)+	31.8	36.9	70.6	74.8	45.5	163.3	11-Jun	44.9	98.2	15.1
Overland +	17.2	25.8	65.2	70.5	43.2	164.7	13-Jun	44.5	100.0	15.4
Peregrine ++	54.0	66.8	96.2	<b>95.2*</b>	<b>58.4*</b>	171.3	19-Jun	48.4	97.1	11.9
R Promontory <sup>1/</sup>	<b>108.3**</b>	<b>117.1**</b>	<b>129.9**</b>	<b>118.0**</b>	<b>59.9**</b>	165.7	14-Jun	45.3	<b>46.3*</b>	11.7
Pryor (P)+	35.3	45.7	68.2	73.9	57.1	172.0	20-Jun	36.5	98.3	13.7
Radiant (P)	86.1	92.2	<b>110.3*</b>	<b>102.6*</b>	<b>59.4*</b>	170.0	18-Jun	47.5	<b>36.1**</b>	12.6
Rampart	48.6	58.1	83.8	82.2	<b>59.0*</b>	170.7	19-Jun	43.2	94.5	14.5
Robidoux +	49.8	59.5	89.7	89.2	48.6	164.0	12-Jun	42.1	91.3	13.7
SY Clearstone 2CL (P, C)	76.2				56.0	168.0	16-Jun	43.8	66.9	13.6
SY Wolf (P)+	60.2	70.4			53.2	163.7	12-Jun	42.6	57.2	15.1
R WB-Quake (P)++	71.8	85.1	<b>110.6*</b>		<b>57.4*</b>	170.6	19-Jun	41.4	82.5	13.9
R Yellowstone +	84.5	96.8	<b>118.6*</b>	<b>113.8*</b>	57.0	170.0	18-Jun	43.4	57.4	13.2
Average	<b>53.1</b>	<b>60.7</b>	<b>85.8</b>	<b>84.5</b>	<b>53.7</b>	<b>168.2</b>	<b>16-Jun</b>	<b>42.1</b>	<b>78.9</b>	<b>14.0</b>
LSD (0.05)	<b>14.6</b>	<b>18.2</b>	<b>23.6</b>	<b>23.4</b>	<b>2.7</b>	<b>2.4</b>		<b>2.3</b>	<b>17.7</b>	
C.V.	<b>16.2</b>	<b>14.7</b>	<b>16.8</b>	<b>19.6</b>	<b>3.0</b>	<b>0.9</b>		<b>3.2</b>	<b>13.5</b>	

\*\* = indicates highest yielding variety within a column CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 4. HARD WINTER : District 2-- Bozeman - Dryland (Moderate Rainfall)

*** No harvest in 2010 due to severe hail ***								
Cultivar/Line	Grain Yield (bushels/acre)			2012 Data				
	2012	2011-12	2009//12	Test weight	Heading Date		Plant height	Protein
					Ordinal	Calendar		
		2 yr	3yr	lb/bu	from Jan1		in	%
Accipiter ++	71.8	66.1	74.3	57.5	171.0	19-Jun	36.8	15.6
AP 503 CL2 (P, CL)+	66.8	63.8	72.3	60.5	165.9	14-Jun	34.2	15.3
Art (P)+	<b>85.5*</b>	78.9		<b>61.0*</b>	164.4	12-Jun	35.4	14.7
Bearpaw	74.2	68.7	76.9	59.2	168.7	17-Jun	35.1	15.4
Broadview	64.1	60.1		57.4	168.8	17-Jun	34.6	15.6
Bynum (P, CL)+	71.3	69.9	73.1	<b>61.3*</b>	167.3	15-Jun	41.2	15.9
R Carter (P)+	70.7	53.8	71.3	59.2	167.6	16-Jun	31.6	15.8
R CDC Falcon (P)+	70.8	63.0	71.1	57.8	168.8	17-Jun	33.3	14.8
Curlew <sup>1/</sup>	76.6	81.8	89.1	57.9	168.7	17-Jun	40.1	14.9
Decade +	78.5	69.1	78.1	59.0	168.0	16-Jun	35.1	15.7
Genou +	68.1	58.0	70.9	59.6	169.7	18-Jun	39.8	16.2
R Jagalene (P)+	<b>87.9*</b>	78.2	81.8	<b>61.3*</b>	166.2	14-Jun	35.3	14.8
Jerry	75.6	71.2	78.3	58.4	170.7	19-Jun	41.8	14.9
Judee ++	71.8	80.5	88.3	58.3	169.3	17-Jun	34.6	16.2
R Ledger (P)+	74.1	60.8	70.8	60.0	167.9	16-Jun	35.7	13.8
McGill +	75.3	64.5		57.4	165.4	13-Jun	37.2	15.2
R MT08172 (Colter)++	75.0	85.6		58.0	170.3	18-Jun	36.2	15.4
MT0871	72.7	67.9		56.2	171.2	19-Jun	34.5	15.8
MT0978	78.6	83.2		57.7	171.0	19-Jun	34.8	15.1
MT10113	81.8			59.2	167.8	16-Jun	36.0	15.3
MT10116	75.3			58.0	170.6	19-Jun	34.7	15.5
MT1078	<b>90.9**</b>			57.9	169.2	17-Jun	33.8	14.2
MT1088	73.3			58.2	170.0	18-Jun	35.9	15.5
MT1090	77.6			56.8	170.5	19-Jun	36.5	15.0
MT1091	83.2			56.5	170.4	18-Jun	35.7	15.2
MT1092	81.7			58.3	170.6	19-Jun	35.9	15.1
MT1105	78.5			57.3	170.1	18-Jun	35.3	15.2
MT1155	80.8			59.3	170.6	19-Jun	34.9	14.9
MT1156	<b>86.3*</b>			59.3	171.5	20-Jun	34.4	14.5
MTCL1067	81.8	84.3		57.9	169.1	17-Jun	38.6	15.3
MTS0808 (Warhorse)++	73.3	81.0		58.8	169.7	18-Jun	34.7	15.6
MTS0819	68.8	71.0		56.4	170.3	18-Jun	32.8	15.4
MTS0819-98 (HWW)	71.8			59.0	169.4	17-Jun	32.2	15.1
MTS1024	<b>89.1*</b>			58.0	170.0	18-Jun	32.8	14.1
MTW08168	78.0	82.1		58.2	174.0	22-Jun	36.3	15.5
R Norris (P, CL)+	77.2	68.2	76.5	60.4	167.0	15-Jun	40.8	14.8
Overland +	<b>90.5*</b>	77.8	80.3	<b>61.4*</b>	165.4	13-Jun	36.8	14.0
Peregrine ++	63.7	75.1	79.5	59.5	170.7	19-Jun	42.1	15.0
R Promontory <sup>1/</sup>	75.7	88.1	93.8	60.6	168.2	16-Jun	36.8	14.5
R Pryor (P)+	74.0	59.1	73.5	58.1	170.9	19-Jun	33.2	15.6
Radiant (P)	62.2	72.8	78.2	58.7	171.2	19-Jun	35.7	14.9
Rampart	69.9	72.3	78.0	60.1	169.6	18-Jun	38.8	16.4
Robidoux +	80.9	82.3	87.5	59.5	166.2	14-Jun	35.2	14.5
SY Clearstone 2CL (P, C)	82.7			56.7	170.4	18-Jun	37.7	15.2
SY Wolf (P)+	<b>84.4*</b>	74.8		<b>61.6**</b>	166.6	15-Jun	33.9	14.7
R WB-Quake (P)++	75.7	79.4		59.1	170.7	19-Jun	34.2	15.7
R Yellowstone +	79.3	85.4	95.8	57.6	170.5	19-Jun	36.2	15.4
<b>Average</b>	<b>76.5</b>	<b>72.9</b>	<b>79.1</b>	<b>58.8</b>	<b>169.2</b>	<b>17-Jun</b>	<b>36.1</b>	<b>15.2</b>
<b>LSD (0.05)</b>	<b>7.6</b>	<b>ns</b>	<b>ns</b>	<b>0.8</b>	<b>0.8</b>		<b>1.7</b>	
<b>C.V.</b>	<b>5.6</b>	<b>16.3</b>	<b>13.9</b>	<b>0.8</b>	<b>0.3</b>		<b>2.7</b>	

\*\* = indicates highest yielding variety within a column      CL = CLEARFIELD wheat tolerant to in      **0.8**  
 \* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's prote      **0.3**  
 R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending  
<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 5. HARD WINTER : District 3-- Huntley - Dryland

Cultivar/Line	Grain Yield (bushels/acre)				2012 Data				
	2012	2011-12	2010-12	2009-12	Test weight lb/bu	Heading Date		Plant height in	Protein %
						Ordinal	Calendar		
		2 yr	3 yr	4 yr		from Jan1			
Accipiter ++	59.9	67.1	70.8	76.5	63.8	155.3	3-Jun	32.7	10.1
AP 503 CL2 (P, CL)+	54.3	57.1	67.0	71.9	<b>65.0*</b>	153.7	2-Jun	30.5	10.3
Art (P)+	59.3	<b>74.2*</b>	<b>78.0*</b>		<b>64.6*</b>	149.7	29-May	32.5	10.9
R Bearpaw	54.1	64.5	72.8	75.2	63.3	156.0	4-Jun	30.4	10.9
Broadview	60.5	66.1	69.9		63.1	156.0	4-Jun	31.1	11.1
Bynum (P, CL)+	55.7	64.4	72.6	70.5	63.8	154.7	3-Jun	33.7	11.9
R Carter (P)+	60.4	64.5	67.2	70.2	63.4	155.3	3-Jun	29.1	11.8
R CDC Falcon (P)+	61.1	64.4	71.1	78.0	63.3	156.0	4-Jun	32.0	10.3
Curlew <sup>1/</sup>	69.6	73.2	<b>81.2*</b>	<b>83.5*</b>	63.2	155.0	3-Jun	37.5	11.2
R Decade +	64.7	67.4	75.6	<b>78.6*</b>	63.5	154.7	3-Jun	32.7	11.3
R Genou +	53.7	63.9	67.8	67.2	63.8	156.3	4-Jun	35.3	11.3
R Jagalene (P)+	56.2	60.5	68.5	78.0	<b>65.1*</b>	152.7	1-Jun	32.8	10.5
Jerry	57.9	60.3	65.5	68.8	62.5	157.7	6-Jun	37.8	10.9
R Judee ++	62.5	68.2	74.6	77.3	63.7	155.3	3-Jun	32.7	12.6
Ledger (P)+	54.9	63.6	73.5	76.2	63.3	155.7	4-Jun	32.3	10.4
McGill +	62.5	<b>75.4*</b>			63.6	153.3	1-Jun	35.8	10.6
R MT08172 (Colter)++	67.1	72.6			62.5	157.7	6-Jun	35.7	11.1
MT0871	<b>70.3*</b>	<b>79.0*</b>	<b>81.1*</b>		61.5	160.3	8-Jun	32.4	11.7
MT0978	62.3	72.6			62.6	159.3	7-Jun	32.1	11.2
MT10113	58.9				63.1	158.3	6-Jun	30.9	10.7
MT10116	<b>72.8*</b>				62.6	156.7	5-Jun	33.5	11.5
MT1078	<b>77.8*</b>				62.0	156.0	4-Jun	33.5	10.1
MT1088	68.2				63.0	155.0	3-Jun	34.8	11.2
MT1090	<b>80.2**</b>				62.5	154.7	3-Jun	36.1	11.3
MT1091	64.3				62.1	156.3	4-Jun	34.0	10.6
MT1092	68.0				63.3	157.3	5-Jun	35.2	10.9
MT1105	<b>72.1*</b>				62.7	155.7	4-Jun	33.7	10.6
MT1155	<b>74.7*</b>				62.2	155.0	3-Jun	34.8	10.6
MT1156	68.7				62.3	158.3	6-Jun	33.1	11.5
MTCL1067	68.4	71.9			62.7	155.7	4-Jun	37.6	10.8
R MTS0808 (Warhorse)++	65.4	72.6	75.0		62.9	158.3	6-Jun	32.7	11.8
MTS0819	63.1	<b>73.6*</b>	<b>78.1*</b>		63.0	156.3	4-Jun	31.9	10.1
MTS0819-98 (HWW)	57.5				64.2	156.7	5-Jun	30.9	11.2
MTS1024	<b>74.4*</b>				61.4	158.7	7-Jun	32.2	9.7
MTW08168	60.9	72.1			62.0	160.7	9-Jun	37.0	11.1
R Norris (P, CL)+	66.2	70.3	<b>77.7*</b>	<b>82.9*</b>	<b>64.4*</b>	152.0	31-May	36.8	11.0
Overland +	<b>73.0*</b>	<b>82.7**</b>	<b>85.3**</b>	<b>87.6**</b>	63.9	151.7	31-May	35.2	12.2
Peregrine ++	65.7	69.3	71.7	74.9	63.0	156.3	4-Jun	37.0	10.5
R Promontory <sup>1/</sup>	70.1	<b>73.4*</b>	<b>76.9*</b>	<b>79.7*</b>	<b>65.2**</b>	154.3	2-Jun	35.5	11.4
R Pryor (P)+	63.6	67.8	71.4	<b>78.1*</b>	62.8	157.3	5-Jun	30.4	10.0
Radiant (P)	56.5	64.0	68.8	73.6	62.3	157.3	5-Jun	34.0	10.7
R Rampart	55.8	64.8	66.7	69.3	63.2	156.0	4-Jun	36.0	12.5
Robidoux +	<b>71.5*</b>	<b>73.5*</b>	<b>78.5*</b>	<b>85.2*</b>	63.9	150.7	30-May	31.8	10.6
SY Clearstone 2CL (P, C)	<b>70.3*</b>				61.8	156.3	4-Jun	35.7	10.8
SY Wolf (P)+	68.7	<b>74.5*</b>			62.9	153.0	1-Jun	32.3	10.9
R WB-Quake (P)++	62.7	71.4	72.0		63.7	158.3	6-Jun	33.4	11.6
R Yellowstone +	69.4	<b>74.9*</b>	<b>79.4*</b>	<b>86.2*</b>	62.4	156.3	4-Jun	34.6	11.0
Average	<b>64.8</b>	<b>69.3</b>	<b>73.5</b>	<b>76.8</b>	<b>63.2</b>	<b>155.8</b>	<b>4-Jun</b>	<b>33.7</b>	<b>11.0</b>
LSD (0.05)	<b>9.0</b>	<b>9.6</b>	<b>9.0</b>	<b>9.5</b>	<b>0.8</b>	<b>2.8</b>		<b>2.2</b>	
C.V.	<b>8.0</b>	<b>6.8</b>	<b>7.5</b>	<b>8.8</b>	<b>0.7</b>	<b>1.1</b>		<b>3.9</b>	

\*\* = indicates highest yielding variety within a column CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 6. HARD WINTER : District 4-- Moccasin - Dryland

Cultivar/Line	Grain Yield (bushels/acre)				2012 Data				
	2012	2011-12	2010-12	2009-12	Test weight lb/bu	Heading Date		Plant height in	Protein %
						Ordinal	Calendar		
		2 yr	3 yr	4 yr		from Jan1			
Accipiter ++	31.0	<b>41.1*</b>	44.8	45.0	57.7	170.0	18-Jun	25.0	13.7
AP 503 CL2 (P, CL)+	28.1	32.3	39.4	40.4	<b>62.9**</b>	166.0	14-Jun	22.1	13.9
Art (P)+	32.4	39.4	<b>48.8*</b>		<b>61.8*</b>	162.3	10-Jun	24.1	13.5
R Bearpaw	32.0	37.8	44.5	43.8	59.3	166.0	14-Jun	23.9	13.3
Broadview	32.9	<b>39.9*</b>	<b>46.9*</b>		58.8	168.7	17-Jun	23.9	14.2
Bynum (P, CL)+	26.2	30.5	35.2	35.4	59.1	166.0	14-Jun	26.0	15.4
R Carter (P)+	<b>35.3*</b>	<b>39.7*</b>	<b>47.3*</b>	46.5	60.5	166.0	14-Jun	24.7	13.3
R CDC Falcon (P)+	<b>34.6*</b>	<b>40.5*</b>	46.4	46.1	57.7	166.7	15-Jun	25.9	13.7
Curlew <sup>1/</sup>	29.8	35.5	42.5	42.9	58.0	167.3	15-Jun	24.9	13.8
R Decade +	31.9	37.4	45.1	45.5	<b>61.6*</b>	166.0	14-Jun	24.5	13.5
R Genou +	28.1	34.0	41.3	41.6	56.3	167.3	15-Jun	26.4	15.4
R Jagalene (P)+	32.6	36.3	45.2	44.5	<b>62.8*</b>	166.0	14-Jun	24.9	12.9
Jerry	29.1	38.3	45.9	45.5	58.5	170.0	18-Jun	25.6	14.4
R Judee ++	30.4	37.0	42.7	41.6	56.4	165.3	13-Jun	25.5	14.9
R Ledger (P)+	28.7	36.0	41.2	41.1	59.3	166.0	14-Jun	24.1	13.6
McGill +	31.5	<b>40.3*</b>			59.2	165.0	13-Jun	24.7	13.4
R MT08172 (Colter)++	<b>36.3*</b>	<b>44.9*</b>			60.0	168.7	17-Jun	24.6	13.4
MT0871	33.8	<b>41.0*</b>	<b>47.7*</b>		57.1	168.7	17-Jun	24.9	15.4
MT0978	<b>34.7*</b>	<b>43.2*</b>			58.2	170.0	18-Jun	24.8	14.3
MT10113	<b>34.8*</b>				59.9	166.0	14-Jun	24.4	13.6
MT10116	34.0				58.6	171.3	19-Jun	24.4	13.8
MT1078	<b>36.1*</b>				58.8	168.7	17-Jun	24.0	13.2
MT1088	<b>36.4*</b>				60.1	169.3	17-Jun	23.1	13.8
MT1090	<b>36.3*</b>				57.2	170.0	18-Jun	27.0	13.2
MT1091	<b>37.6*</b>				58.6	168.7	17-Jun	27.1	13.2
MT1092	<b>37.7**</b>				59.9	169.3	17-Jun	26.5	13.6
MT1105	<b>36.1*</b>				59.0	170.0	18-Jun	26.0	13.3
MT1155	32.8				59.2	170.0	18-Jun	24.1	13.8
MT1156	34.1				59.5	170.0	18-Jun	24.0	14.0
MTCL1067	32.6	<b>42.5*</b>			60.0	168.7	17-Jun	26.1	13.5
R MTS0808 (Warhorse)++	31.5	39.4	45.8		60.0	170.0	18-Jun	21.4	13.4
MTS0819	31.4	<b>39.9*</b>	46.6		57.8	168.7	17-Jun	22.9	13.3
MTS0819-98 (HWW)	30.2				58.9	170.0	18-Jun	23.2	13.8
MTS1024	<b>35.3*</b>				57.3	168.7	17-Jun	25.2	13.0
MTW08168	33.5	<b>42.1*</b>			59.2	170.0	18-Jun	24.7	14.4
R Norris (P, CL)+	31.3	36.5	43.1	43.3	59.3	171.0	19-Jun	25.3	14.2
Overland +	34.2	<b>43.9*</b>	<b>51.8*</b>	<b>49.5*</b>	60.2	166.0	14-Jun	25.5	13.6
Peregrine ++	28.7	35.1	40.0	40.2	57.7	164.0	12-Jun	26.5	13.9
R Promontory <sup>1/</sup>	28.7	36.7	44.3	44.7	59.3	168.7	17-Jun	24.2	14.2
R Pryor (P)+	<b>37.7**</b>	<b>41.6*</b>	<b>48.5*</b>	<b>48.3*</b>	59.4	167.3	15-Jun	24.0	12.3
Radiant (P)	27.1	34.1	35.9	36.1	58.3	168.7	17-Jun	24.5	14.2
R Rampart	28.5	33.0	38.2	38.7	59.0	170.0	18-Jun	25.7	15.1
Robidoux +	<b>34.4*</b>	<b>45.1**</b>	<b>51.3*</b>	<b>50.5*</b>	58.6	167.3	15-Jun	25.5	14.1
SY Clearstone 2CL (P, C)	<b>34.8*</b>				57.8	165.7	14-Jun	25.5	13.4
SY Wolf (P)+	33.5	<b>39.9*</b>			61.1	166.0	14-Jun	24.0	13.5
R WB-Quake (P)++	30.0	<b>40.0*</b>	43.3		57.3	170.0	18-Jun	23.8	14.7
R Yellowstone +	35.5*	<b>43.7*</b>	<b>52.5**</b>	<b>52.2**</b>	58.3	170.0	18-Jun	26.3	14.0
Average	<b>32.7</b>	<b>38.8</b>	<b>44.5</b>	<b>43.8</b>	<b>59.1</b>	<b>168.0</b>	<b>16-Jun</b>	<b>24.8</b>	<b>13.8</b>
LSD (0.05)	<b>3.5</b>	<b>15.7</b>	<b>5.8</b>	<b>4.6</b>	<b>1.4</b>	<b>2.7</b>		<b>2.5</b>	
C.V.	<b>6.0</b>	<b>7.2</b>	<b>8.0</b>	<b>7.4</b>	<b>1.2</b>	<b>1.0</b>		<b>5.7</b>	

\*\* = indicates highest yielding variety within a column

CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 7. HARD WINTER : District 5-- Conrad - Dryland

Cultivar/Line	Grain Yield (bushels/acre)				2012 Data				
	2012	2011-12	2010-12	2009-12	Test weight lb/bu	Heading Date		Plant height in	Protein %
						Ordinal	Calendar		
		2 yr	3 yr	4 yr		from Jan1			
Accipiter ++	96.9*	96.1*	98.6*	92.1*	62.1*	168.2	16-Jun	33.4	10.6
AP 503 CL2 (P, CL)+	70.5	76.5	79.0	79.0	63.1**	166.6	15-Jun	27.9	12.2
Art (P)+	85.4	94.1*	89.9		61.9*	164.0	12-Jun	30.4	12.2
R Bearpaw	78.1	84.3	89.7	85.0	61.4	167.3	15-Jun	30.8	12.0
Broadview	87.7*	92.3*	99.0*		61.2	167.8	16-Jun	31.5	11.4
R Bynum (P, CL)+	66.6	69.7	76.2	73.7	60.8	165.9	14-Jun	32.4	13.0
R Carter (P)+	85.1	88.2	87.5	83.2	61.4	167.1	15-Jun	29.8	11.8
R CDC Falcon (P)+	92.9*	91.1*	96.3*	93.4*	61.8*	166.8	15-Jun	30.1	11.3
Curlew <sup>1/</sup>	83.7	89.2	92.8	90.5*	60.9	167.0	15-Jun	35.2	11.9
R Decade +	81.5	86.6	90.6	87.7	62.4*	166.3	14-Jun	29.8	12.7
R Genou +	77.3	80.7	85.5	83.7	61.6	167.7	16-Jun	34.4	12.0
R Jagalene (P)+	92.9*	94.9*	94.8*	89.8	62.9*	167.0	15-Jun	32.9	12.3
Jerry	78.6	81.8	83.2	80.9	60.5	168.2	16-Jun	36.5	11.6
R Judee ++	95.4*	96.4*	96.3*	90.6*	63.0*	168.3	16-Jun	32.3	11.8
R Ledger (P)+	78.2	83.8	87.9	84.7	62.3*	167.5	16-Jun	31.4	11.2
McGill +	84.6	93.1*			60.9	164.4	12-Jun	33.3	12.0
R MT08172 (Colter)++	95.3*	97.5*			61.1	168.9	17-Jun	33.5	11.6
MT0871	89.3*	97.4*	101.5*		60.2	171.2	19-Jun	32.2	11.8
MT0978	97.0*	98.2*			60.7	170.6	19-Jun	32.3	11.6
MT10113	88.1*				62.0*	165.9	14-Jun	31.7	12.3
MT10116	95.2*				60.8	169.3	17-Jun	32.9	12.1
MT1078	95.8*				59.9	168.8	17-Jun	31.3	11.3
MT1088	88.7*				60.2	168.3	16-Jun	33.5	11.8
MT1090	101.1**				61.0	167.8	16-Jun	34.6	11.3
MT1091	92.6*				60.2	167.9	16-Jun	33.2	11.3
MT1092	89.7*				60.5	169.5	18-Jun	33.8	11.8
MT1105	90.7*				60.4	168.5	17-Jun	30.6	11.7
MT1155	87.1*				59.1	168.7	17-Jun	33.2	11.4
MT1156	87.3*				60.9	169.4	17-Jun	32.6	11.3
MTCL1067	84.0	89.8			60.9	168.4	16-Jun	34.1	11.9
R MTS0808 (Warhorse)++	84.5	89.4	93.9*		61.0	168.2	16-Jun	31.9	11.6
MTS0819	88.6*	95.4*	101.6*		60.4	169.0	17-Jun	31.5	11.9
MTS0819-98 (HWW)	91.7*				62.4*	168.3	16-Jun	30.2	11.6
MTS1024	93.9*				60.3	169.0	17-Jun	30.7	11.2
MTW08168	88.8*	96.4*			61.3	173.3	21-Jun	33.6	12.1
Norris (P, CL)+	85.0	87.1	89.9	85.2	62.4*	166.0	14-Jun	33.1	11.7
Overland +	87.0*	93.0*	99.5*	94.0*	62.1*	166.0	14-Jun	31.9	12.2
Peregrine ++	79.5	82.4	86.0	85.5	60.6	167.1	15-Jun	37.7	10.9
Promontory <sup>1/</sup>	79.8	89.7	94.0*	89.7	62.9*	167.0	15-Jun	32.2	11.2
R Pryor (P)+	92.5*	97.3*	98.9*	95.1*	60.6	169.2	17-Jun	31.5	11.4
Radiant (P)	81.2	84.4	89.1	86.4	60.0	168.0	16-Jun	35.0	12.1
R Rampart	73.4	74.9	77.4	76.3	60.3	168.6	17-Jun	33.3	12.8
Robidoux +	94.2*	97.9*	98.7*	96.8*	62.4*	165.8	14-Jun	31.6	11.2
SY Clearstone 2CL (P, C)	86.8*				60.2	169.2	17-Jun	33.1	11.9
SY Wolf (P)+	88.3*	99.0**			61.9*	166.0	14-Jun	29.2	11.5
R WB-Quake (P)++	87.7*	87.4	90.0		61.5	169.2	17-Jun	32.0	11.7
R Yellowstone +	96.2*	98.6*	102.0**	97.3**	60.6	168.6	17-Jun	32.4	11.1
Average	87.2	89.8	91.8	87.3	61.2	167.9	16-Jun	32.4	11.7
LSD (0.05)	15.4	9.6	8.4	7.0	1.3	1.4		2.4	
C.V.	10.1	4.7	5.6	5.7	1.2	0.5		4.4	

\*\* = indicates highest yielding variety within a column

CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 8. HARD WINTER : District 5-- Havre - Dryland

*** No harvest in 2011 due to uneven stands ***									
Cultivar/Line	Grain Yield (bushels/acre)			2012 Data					
	2012	2010//12	2009//12	Test weight	Heading Date		Plant height <sup>2/</sup>	Sawfly cutting <sup>2/</sup>	Protein
					Ordinal <sup>2/</sup>	Calendar			
		2 yr	3 yr	lb/bu	from Jan1		in	%	%
Accipiter ++	47.5	60.6	54.9	57.8	167.4	15-Jun	29.5	10	13.2
AP 503 CL2 (P, CL)+	54.0	<b>64.4*</b>	53.4	<b>60.1*</b>	160.9	9-Mar	29.1	11	12.9
Art (P)+	56.4	<b>68.8**</b>		<b>60.7*</b>	158.3	6-Jun	29.9	7	12.5
R Bearpaw	55.4	<b>61.7*</b>	53.7	58.8	164.9	13-Jun	30.2	3	13.1
Broadview	52.8	<b>63.2*</b>		58.2	164.3	12-Jun	32.1	9	11.9
R Bynum (P, CL)+	43.4	53.0	48.5	59.3	164.1	12-Jun	32.7	3	13.8
R Carter (P)+	52.1	<b>63.4*</b>	54.8	58.8	167.5	16-Jun	28.1	1	13.5
R CDC Falcon (P)+	53.3	<b>62.8*</b>	55.4	59.3	164.7	13-Jun	28.4	7	13.2
Curlew <sup>1/</sup>	44.6	53.3	47.3	56.8	165.3	13-Jun	33.5	14	13.9
R Decade +	55.8	<b>64.0*</b>	54.5	59.7	168.5	17-Jun	30.9	10	12.6
R Genou +	43.5	52.6	50.1	58.1	166.8	15-Jun	34.1	6	14.3
R Jagalene (P)+	53.8	<b>65.8*</b>	54.9	<b>61.6**</b>	159.7	8-Jun	29.5	9	13.3
Jerry	44.2	56.4	49.6	57.7	166.0	14-Jun	33.8	14	13.2
R Judee ++	48.9	60.7	54.8	56.9	167.2	15-Jun	30.5	2	14.6
R Ledger (P)+	48.1	59.0	52.7	<b>60.0*</b>	166.6	15-Jun	27.4	5	12.2
McGill +	56.9			59.3	159.3	7-Jun	31.0	13	11.8
R MT08172 (Colter)++	<b>70.7**</b>			58.7	167.2	15-Jun	31.1	11	11.9
MT0871	58.4	<b>67.6*</b>		57.8	168.3	16-Jun	29.9	12	14.1
MT0978	54.8			58.5	167.0	15-Jun	30.2	9	13.3
MT10113	58.0			59.6	162.6	11-Jun	29.4	6	13.3
MT10116	54.5			58.0	167.5	16-Jun	30.7	11	14.4
MT1078	<b>62.0*</b>			57.0	165.6	14-Jun	29.1	4	12.0
MT1088	46.9			58.3	166.3	14-Jun	30.8	20	14.2
MT1090	58.3			56.3	166.4	14-Jun	31.7	11	13.4
MT1091	56.3			57.2	166.8	15-Jun	30.6	10	13.4
MT1092	51.9			57.7	165.8	14-Jun	29.4	11	12.9
MT1105	<b>61.1*</b>			58.1	166.1	14-Jun	30.1	11	12.5
MT1155	49.6			57.8	167.0	15-Jun	31.8	14	14.3
MT1156	53.1			57.4	168.0	16-Jun	30.6	11	14.6
MTCL1067	57.0			58.2	165.0	13-Jun	33.1	7	13.5
R MTS0808 (Warhorse)++	51.7	<b>61.7*</b>		57.6	166.0	14-Jun	30.0	2	13.1
MTS0819	46.7	<b>62.0*</b>		55.8	168.0	16-Jun	30.4	6	14.5
MTS0819-98 (HWW)	54.0			58.4	167.3	15-Jun	27.4	3	12.7
MTS1024	55.3			57.2	167.0	15-Jun	30.7	6	12.6
MTW08168	52.1			59.0	168.7	17-Jun	35.2	18	13.1
Norris (P, CL)+	55.0	60.4	53.1	59.4	158.9	7-Jun	33.4	10	12.5
Overland +	49.0	<b>65.4*</b>	57.4	59.0	164.0	12-Jun	28.9	3	13.1
Peregrine ++	43.9	55.8	49.4	56.9	166.3	14-Jun	39.4	16	13.0
Promontory <sup>1/</sup>	45.4	<b>61.9*</b>	53.4	<b>61.2*</b>	162.6	11-Jun	30.5	16	12.4
R Pryor (P)+	57.6	<b>65.6*</b>	57.1	58.0	167.0	15-Jun	30.1	3	12.2
Radiant (P)	47.2	55.9	50.3	57.5	166.7	15-Jun	33.2	15	14.3
R Rampart	46.6	52.7	48.2	58.7	166.6	15-Jun	31.1	0	13.5
Robidoux +	54.5	<b>62.9*</b>	53.8	59.1	161.8	10-Jun	29.8	7	13.3
SY Clearstone 2CL (P, C)	59.6			57.1	165.9	14-Jun	32.2	11	13.3
SY Wolf (P)+	56.2			<b>60.7*</b>	161.2	9-Jun	31.4	5	13.3
R WB-Quake (P)++	48.4	59.5		58.8	169.9	18-Jun	30.2	4	14.3
R Yellowstone +	52.1	<b>62.7*</b>	55.3	58.1	166.3	14-Jun	31.7	9	13.5
<b>Average</b>	<b>52.9</b>	<b>60.8</b>	<b>52.8</b>	<b>58.5</b>	<b>165.4</b>	<b>13-Jun</b>	<b>31.0</b>	<b>8.4</b>	<b>13.2</b>
<b>LSD (0.05)</b>	<b>9.7</b>	<b>8.0</b>	<b>ns</b>	<b>1.7</b>	<b>3.2</b>		<b>2.4</b>	<b>8.4</b>	
<b>C.V.</b>	<b>10.5</b>	<b>6.4</b>	<b>8.7</b>	<b>1.6</b>	<b>1.1</b>		<b>4.5</b>	<b>7.1</b>	

\*\* = indicates highest yielding variety within a column

CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat<sup>2/</sup> = 2011 data



Table 9. HARD WINTER : District 6-- Sidney - Dryland

*** No harvest in 2009 due to severe winter-kill ***								
Cultivar/Line	Grain Yield (bushels/acre)			Test weight	Heading Date		Plant height	Protein %
	2012	2011-12	2010-12		Ordinal	Calendar		
		2 yr	3 yr	lb/bu	from Jan1		in	%
Accipiter ++	59.4*	58.9*	62.7*	56.0	159.5	8-Jun	29.7	11.6
AP 503 CL2 (P, CL)+	55.3	48.5	47.9	60.0	155.5	4-Jun	26.6	11.1
Art (P)+	46.7	49.2	48.7	60.5	155.5	4-Jun	26.6	12.5
Bearpaw	60.3*	57.5*	54.8	58.0	156.5	5-Jun	28.0	11.5
Broadview	58.1*	56.8	60.4*	57.5	157.0	5-Jun	28.5	11.3
Bynum (P, CL)+	48.3	42.0	42.2	59.5	156.5	5-Jun	30.7	12.9
R Carter (P)+	49.8	51.3	50.6	59.5	158.0	6-Jun	26.2	11.9
R CDC Falcon (P)+	61.9*	59.4*	59.9*	58.5	155.5	4-Jun	27.8	12.8
Curlew <sup>1/</sup>	58.4*	56.5	54.3	59.0	157.0	5-Jun	32.7	12.2
R Decade +	55.3	60.2*	60.2*	60.0	157.0	5-Jun	28.0	12.2
Genou +	53.4	48.5	48.6	59.0	156.5	5-Jun	33.3	14.0
Jagalene (P)+	57.3*	57.5*	54.4	60.5	155.5	4-Jun	28.0	11.9
R Jerry	65.9*	67.1*	66.1*	58.0	157.0	5-Jun	33.7	11.8
Judee ++	51.9	48.2	49.4	58.0	158.0	6-Jun	28.5	11.4
Ledger (P)+	46.3	45.3	45.9	58.5	158.0	6-Jun	28.3	11.8
McGill +	53.5	55.0		58.5	156.0	4-Jun	29.7	11.9
MT08172 (Colter)++	62.5*	64.5*		59.0	160.0	8-Jun	30.9	10.9
MT0871	63.7*	60.6*	61.4*	58.5	159.0	7-Jun	28.7	11.3
MT0978	63.5*	65.9*		60.0	158.0	6-Jun	29.3	11.9
MT10113	53.2			58.5	156.5	5-Jun	27.4	11.4
MT10116	56.3			59.0	159.0	7-Jun	28.1	11.0
MT1078	60.1*			56.5	157.5	6-Jun	29.7	10.1
MT1088	59.6*			57.5	158.5	7-Jun	31.3	12.0
MT1090	66.0*			57.0	156.5	5-Jun	30.7	10.9
MT1091	64.0*			55.5	158.0	6-Jun	29.3	11.8
MT1092	60.4*			58.0	158.5	7-Jun	29.9	11.4
MT1105	62.0*			57.0	157.5	6-Jun	30.1	11.7
MT1155	55.0			55.0	159.0	7-Jun	29.7	11.7
MT1156	54.4			58.5	161.0	9-Jun	29.9	11.7
MTCL1067	59.8*	58.1*		57.5	157.5	6-Jun	33.3	10.6
MTS0808 (Warhorse)++	50.9	55.2	55.2	56.0	158.0	6-Jun	28.0	11.4
MTS0819	54.3	52.0	54.4	59.0	160.0	8-Jun	27.0	12.1
MTS0819-98 (HWW)	52.6			61.0	157.5	6-Jun	28.0	11.0
MTS1024	60.1*			57.5	158.0	6-Jun	30.7	11.9
MTW08168	59.7*	62.0*		58.0	160.0	8-Jun	33.1	11.1
Norris (P, CL)+	54.7	57.2*	56.8	60.5	155.0	3-Jun	30.7	11.9
Overland +	56.8	67.4**	66.7**	60.5	156.0	4-Jun	28.7	11.7
Peregrine ++	54.4	55.3	55.5	57.5	159.0	7-Jun	36.4	12.7
Promontory <sup>1/</sup>	64.3*	57.6*	57.8	60.5	156.5	5-Jun	29.7	11.9
R Pryor (P)+	62.4*	60.0*	55.2	57.5	158.0	6-Jun	28.9	10.6
Radiant (P)	55.9	54.6	55.2	56.0	157.5	6-Jun	31.3	11.0
Rampart	43.9	45.7	44.1	57.5	159.0	7-Jun	31.1	11.8
Robidoux +	56.4	52.6	53.2	58.5	156.0	4-Jun	26.6	12.1
SY Clearstone 2CL (P, C)	62.9*			57.0	158.0	6-Jun	29.9	11.0
SY Wolf (P)+	54.1	56.5		60.5	156.0	4-Jun	28.0	11.2
R WB-Quake (P)++	49.4	51.1	50.6	56.5	159.0	7-Jun	29.3	13.5
Yellowstone +	63.7*	65.6*	64.4*	59.5	157.0	5-Jun	31.3	12.4
Average	57.4	56.0	54.9	58.4	157.5	6-Jun	29.7	11.7
LSD (0.05)	9.6	10.4	8.1		1.9		2.6	
C.V.	8.3	9.1	9.0		0.6		4.4	

\*\* = indicates highest yielding variety within a column

CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 10. HARD WINTER : District 6-- Williston, North Dakota - Dryland

*** No harvest in 2010 due to severe winterkill ***										
Cultivar/Line	Grain Yield (bushels/acre)			2012 Data						
	2012	2011-12	2009//12	Test weight	Winter survival	Heading Date		Plant height	1000 kernel wt	Protein
	2 yr		3 yr	lb/bu	%	Ordinal	Calendar	in	g	%
Accipiter ++	46.7	56.2	52.5*	58.3*	48.3*	155.5	4-Jun	27.2	25.4	12.5
AP 503 CL2 (P, CL)+	33.4	45.9	43.7	59.2*	13.3	155.0	3-Jun	26.4	30.1	14.3
Art (P)+	29.6	43.0		57.8	8.3	150.0	30-May	26.5	27.1	14.5
Bearpaw	46.6	56.0	49.4*	57.5	25.0	154.5	3-Jun	25.3	27.2	13.5
Broadview	62.6**	64.6		58.9*	51.7*	154.0	2-Jun	26.8	28.9	12.6
Bynum (P, CL)+	33.3	42.6	41.0	57.3	11.7	155.0	3-Jun	28.5	29.0	15.1
R Carter (P)+	41.9	50.6	48.8*	58.4*	18.3	156.0	4-Jun	25.6	25.2	14.6
R CDC Falcon (P)+	59.6*	63.1	58.2**	59.2*	51.7*	153.0	1-Jun	26.8	28.1	11.9
Curlew <sup>1/</sup>	25.3	43.5	42.5	53.4	8.3	156.5	5-Jun	31.0	24.9	15.8
R Decade +	43.4	56.3	50.7*	58.7*	46.7*	153.0	1-Jun	27.6	30.3	14.1
Genou +	29.3	41.4	41.5	55.9	10.3	157.0	5-Jun	26.8	24.7	14.7
Jagalene (P)+	38.1	46.3	44.8	59.4**	11.7	153.0	1-Jun	25.7	31.2	13.8
R Jerry	53.7*	54.2	52.4*	58.0	36.7	154.5	3-Jun	29.8	32.8	13.1
Judee ++	21.5	43.3	42.1	55.0	5.3	158.0	6-Jun	27.0	27.8	16.1
Ledger (P)+	34.6	45.4	45.2	57.9	21.7	155.5	4-Jun	27.8	31.9	13.5
McGill +	43.7	53.5		56.9	25.0	152.5	1-Jun	28.2	30.1	14.1
MT08172 (Colter)++	41.7	52.4		57.2	18.3	156.5	5-Jun	27.2	32.8	14.3
MT0871	48.0	53.9		56.0	20.0	158.0	6-Jun	26.5	25.2	13.8
MT0978	41.2	54.0		56.7	13.3	156.5	5-Jun	27.2	27.7	14.5
MT10113	40.0			58.5*	28.3	154.0	2-Jun	25.2	28.0	13.6
MT10116	44.6			57.3	18.3	156.0	4-Jun	27.5	29.7	13.8
MT1078	43.6			55.5	15.0	157.0	5-Jun	27.6	27.5	13.5
MT1088	43.6			57.5	25.0	156.0	4-Jun	28.7	30.6	13.6
MT1090	57.9*			57.9	43.3*	155.5	4-Jun	29.1	31.9	13.0
MT1091	42.3			55.6	18.3	157.0	5-Jun	29.4	29.3	13.3
MT1092	38.5			55.6	11.7	158.0	6-Jun	26.9	29.8	14.4
MT1105	42.4			56.2	18.3	156.0	4-Jun	26.3	29.6	13.6
MT1155	40.3			55.3	23.3	156.5	5-Jun	28.8	23.5	14.3
MT1156	51.2*			57.6	31.7	156.0	4-Jun	28.6	28.8	13.1
MTCL1067	32.2	48.5		55.7	8.3	156.0	4-Jun	28.6	31.4	15.2
MTS0808 (Warhorse)++	46.7	51.2		56.6	23.3	156.5	5-Jun	26.7	26.9	14.1
MTS0819	42.1	51.2		57.1	25.0	155.5	4-Jun	27.8	26.7	14.7
MTS0819-98 (HWW)	27.0			56.9	12.0	156.5	5-Jun	26.5	26.2	14.7
MTS1024	34.7			55.0	13.3	156.5	5-Jun	27.8	27.8	14.5
MTW08168	52.9*	54.6		57.9	26.7	157.5	6-Jun	29.3	28.5	13.3
Norris (P, CL)+	44.2	49.7	49.6*	59.0*	20.0	151.5	31-May	27.6	32.1	13.9
Overland +	49.4	57.0	51.6*	59.0*	28.3	154.5	3-Jun	26.8	32.3	13.4
Peregrine ++	59.3*	59.6	55.8*	58.2*	56.7**	155.5	4-Jun	31.8	27.5	13.4
Promontory <sup>1/</sup>	30.1	42.7	44.1	57.3	6.0	155.5	4-Jun	27.4	29.1	13.9
R Pryor (P)+	43.9	55.5	51.4*	57.1	16.7	157.0	5-Jun	26.9	25.1	12.7
Radiant (P)	49.5	56.2	54.2*	57.3	28.3	156.0	4-Jun	29.7	30.0	13.1
Rampart	36.0	50.9	47.4	58.1*	18.7	156.5	5-Jun	28.8	24.7	13.9
Robidoux +	40.8	53.0	47.6	56.6	10.0	152.0	31-May	27.1	28.6	14.1
SY Clearstone 2CL (P, C)	38.6			54.9	10.0	156.5	5-Jun	29.8	31.3	14.9
SY Wolf (P)+	27.3	43.3		57.5	6.7	153.0	1-Jun	25.7	31.2	14.0
R WB-Quake (P)++	47.5	52.3		56.7	20.0	157.5	6-Jun	28.1	25.0	14.3
Yellowstone +	43.9	54.9	53.6*	55.8	20.0	156.0	4-Jun	28.7	30.6	14.6
Average	42.2	51.4	48.6	57.2	21.8	155.5	4-Jun	27.7	28.7	13.9
LSD (0.05)	12.9	ns	10.3	1.4	14.7	2.1		3.0		
C.V.	17.6	13.8	12.9	1.2	41.6	0.7		6.6		

\*\* = indicates highest yielding variety within a column CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

R = Recommended Variety; (P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

Table 11. 2005//2012 Intrastate Winter Wheat Test (Exp. 35): Combined Locations Winter Survival and associated Yield (Locations: Williston (2005-2008, 2012), Sidney (2005, 2006, 2008, 2010, 2011) = 10 locations

location-years	*** No recordable Winterkill in 2009 ***															
	Winter Survival (%)							Yield under Winterkill conditions								
	2012	2011-12	2010-12	2008-12	2007-12	2006-12	2005-12	2012	2011-12	2010-12	2008-12	2007-12	2006-12	2005-12		
	1	2	3	5	6	8	10	1	2	3	5	6	8	10		
<b>Accipiter ++</b>	<b>48.3*</b>	<b>60.6*</b>	<b>64.3*</b>	<b>62.2**</b>				46.7	<b>52.5*</b>	<b>58.4*</b>	<b>49.8*</b>					
<b>AP 503 CL2 (P, CL)+</b>	13.3	27.1	32.5	24.7				33.4	37.6	40.6	29.7					
<b>Art (P)+</b>	8.3	17.6	23.9					29.6	40.7	43.0						
<b>Bearpaw</b>	25.0	42.0	43.7					46.6	<b>50.6*</b>	50.2						
<b>Broadview</b>	<b>51.7*</b>	<b>56.4*</b>	<b>61.4*</b>					<b>62.6**</b>	<b>59.1*</b>	<b>61.9*</b>						
<b>Bynum (P, CL)+</b>	11.7	31.2	32.5	22.1	24.3	28.0	33.5	33.3	34.5	37.2	24.9	29.0	30.3	32.3		
<b>Carter (P)+</b>	18.3	30.5	33.6	25.9	31.9	35.1	44.3	41.9	47.3	48.0	36.5	41.4	40.1	43.1		
<b>CDC Falcon (P)+</b>	<b>51.7*</b>	<b>60.0*</b>	<b>62.0*</b>	49.2	<b>53.5*</b>	<b>55.9*</b>	<b>62.0*</b>	<b>59.6*</b>	<b>58.3*</b>	<b>59.1*</b>	<b>49.2*</b>	<b>53.5*</b>	<b>52.4*</b>	<b>53.6*</b>		
<b>Curlew<sup>1/</sup></b>	8.3	29.7	34.4					25.3	39.9	43.2						
<b>Decade +</b>	<b>46.7*</b>	<b>60.9*</b>	59.0	51.8	<b>55.9**</b>			43.4	<b>54.2*</b>	<b>56.2*</b>	<b>46.6*</b>	<b>51.8*</b>				
<b>Genou +</b>	10.3	32.3	37.6	26.7	32.0	36.2	44.2	29.3	36.4	40.6	30.1	34.6	35.8	38.2		
<b>Jagalene (P)+</b>	11.7	32.8	35.4	27.1	31.2	37.4	45.4	38.1	47.9	48.0	37.9	41.7	41.7	44.2		
<b>Jerry</b>	36.7	<b>53.5*</b>	58.7	49.5	<b>55.7*</b>	<b>58.7**</b>	<b>65.0**</b>	<b>53.7*</b>	<b>61.0*</b>	<b>62.1*</b>	<b>51.6**</b>	<b>55.7**</b>	<b>54.4**</b>	<b>56.3**</b>		
<b>Judee ++</b>	5.3	27.7	36.6	27.0				21.5	33.0	39.3	28.9					
<b>Ledger (P)+</b>	21.7	33.4	36.6	26.8	30.7	34.3	41.2	34.6	39.4	42.0	30.5	35.2	36.2	38.3		
<b>McGill +</b>	25.0	<b>55.0*</b>						43.7	<b>50.1*</b>							
<b>MT08172 (Colter)++</b>	18.3	37.2						41.7	<b>54.1*</b>							
<b>MT0871</b>	20.0	<b>46.7*</b>	48.7					48.0	<b>52.7*</b>	<b>56.1*</b>						
<b>MT0978</b>	13.3	40.3						41.2	<b>54.7*</b>							
<b>MT10113</b>	28.3							40.0								
<b>MT10116</b>	18.3							44.6								
<b>MT1078</b>	15.0							43.6								
<b>MT1088</b>	25.0							43.6								
<b>MT1090</b>	<b>43.3*</b>							<b>57.9*</b>								
<b>MT1091</b>	18.3							42.3								
<b>MT1092</b>	11.7							38.5								
<b>MT1105</b>	18.3							42.4								
<b>MT1155</b>	23.3							40.3								
<b>MT1156</b>	31.7							<b>51.2*</b>								
<b>MTCL1067</b>	8.3	24.4						32.2	44.3							
<b>MTS0808 (Warhorse)+</b>	23.3	44.2	44.2					46.7	<b>53.1*</b>	53.8						
<b>MTS0819</b>	25.0	34.3	40.8					42.1	45.9	50.3						
<b>MTS0819-98 (HWW)</b>	12.0							27.0								
<b>MTS1024</b>	13.3							34.7								
<b>MTW08168</b>	26.7	35.0						<b>52.9*</b>	<b>58.6*</b>							
<b>Norris (P, CL)+</b>	20.0	40.5	44.7	33.8	37.6	42.7	49.4	44.2	<b>51.9*</b>	53.3	40.8	44.9	44.6	46.0		
<b>Overland +</b>	28.3	45.3	47.8					49.4	<b>63.7**</b>	<b>64.2**</b>						
<b>Peregrine ++</b>	<b>56.7**</b>	<b>66.4**</b>	<b>70.3**</b>	<b>54.9*</b>				<b>59.3*</b>	<b>57.7*</b>	<b>57.1*</b>	<b>47.7*</b>					
<b>Promontory<sup>1/</sup></b>	6.0	23.6	30.3	24.0	27.5	33.6	38.7	30.1	40.5	46.4	34.0	38.4	39.3	40.6		
<b>Pryor (P)+</b>	16.7	35.2	37.9	32.2	38.5	42.1	50.3	43.9	<b>50.7*</b>	49.0	40.6	44.6	44.7	47.7		
<b>Radiant (P)</b>	28.3	<b>51.2*</b>	54.7					49.5	<b>51.4*</b>	53.0						
<b>Rampart</b>	18.7	34.0	33.2	22.6	25.5	28.5	35.6	36.0	41.7	41.4	28.5	32.2	32.9	35.7		
<b>Robidoux +</b>	10.0	21.9	29.7					40.8	44.8	48.0						
<b>SY Clearstone 2CL (P)</b>	10.0							38.6								
<b>SY Wolf (P)+</b>	6.7	27.5						27.3	43.1							
<b>WB-Quake (P)++</b>	20.0	39.0	42.6					47.5	<b>50.1*</b>	49.9						
<b>Yellowstone +</b>	20.0	36.8	41.3	33.1	37.6	43.2	51.4	43.9	<b>55.7*</b>	<b>57.8*</b>	44.8	49.2	48.8	51.1		
<b>Average</b>	<b>21.8</b>	<b>39.2</b>	<b>43.5</b>	<b>34.9</b>	<b>37.1</b>	<b>39.6</b>	<b>46.8</b>	<b>42.2</b>	<b>48.7</b>	<b>50.4</b>	<b>38.3</b>	<b>42.5</b>	<b>41.8</b>	<b>43.9</b>		
<b>LSD (0.05)</b>	<b>14.7</b>	<b>16.9</b>	<b>11.1</b>	<b>9.5</b>	<b>7.0</b>	<b>5.8</b>	<b>5.9</b>	<b>12.9</b>	<b>13.9</b>	<b>9.5</b>	<b>6.4</b>	<b>4.9</b>	<b>4.2</b>	<b>3.6</b>		
<b>C.V.</b>	<b>41.6</b>	<b>21.2</b>	<b>15.6</b>	<b>21.5</b>	<b>16.4</b>	<b>14.7</b>	<b>14.2</b>	<b>17.6</b>	<b>14.0</b>	<b>11.6</b>	<b>13.3</b>	<b>10.0</b>	<b>10.1</b>	<b>9.2</b>		

\*\* = indicates highest yielding variety within a column

CL = CLEARFIELD wheat tolerant to imidazolinone (IMI) herbicides

\* = indicates varieties yielding equal to highest yielding variety within a column based on Fisher's protected LSD (p=0.05)

(P) = Private Variety; + = Protected Variety; ++ = PVP Pending

<sup>1/</sup> = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat

**Table 12. HARD WINTER WHEAT: Yield Performance under Sawfly Pressure and % Sawfly Cutting (2005-2012) Cutting (Note: Sawfly cutting in each location-year >10%)**

Cultivar/Line	Grain Yield (bu/a)																	Sawfly Cutting (%)																
	2012			2011-12			2010-12			2009-12			2008-12			2007-12			2006-12			2005-12												
	3	5	7	5	7	9	7	9	11	13	15	17	3	5	7	9	11	13	15	17														
Location-years	36.9	42.3	47.4	46.8*	39.8	34.9	40.3	39.1	37.8	34.0	11.8*	9.8*	8.4*	39.2	28.9	35.9	36.7	37.1	37.5	35.9														
Accipiter	33.6	37.9	48.2*	49.8*	36.1	28.7	34.9	33.6	33.9	15.8*	20.5*	21.6	20.1	18.2	20.0	46.1	38.3	43.2	43.9	44.7	42.5	40.6												
AP503 CL2 (P)+	37.3	43.0*	48.2*	49.3*	45.0	37.0	44.2	47.0	44.5	45.0	37.0	44.2	47.0	44.1	44.3	45.0	37.0	44.2	47.0	44.5	46.4	44.3												
Bearpaw ++ ss	35.7	43.8*	49.6*	50.3*	36.7	42.2	46.3	45.8	47.5	49.4*	49.2*	49.3*	49.3*	30.5	26.0	35.3	33.0	32.6	33.7	33.4	43.8	42.0												
CDC Falcon (P)+	39.0	44.4*	50.8*	50.3*	37.0	40.4	44.7	43.3	45.1	46.5	45.8	45.4	45.4	44.0	34.1	44.0	34.1	2.8**	2.9**	9.9*	40.9	8.2*	8.4*											
Decade +	31.4	38.6	46.8	47.7	32.1	40.4	44.7	43.3	45.1	46.5	45.8	45.4	45.4	30.5	26.0	35.3	33.0	32.6	33.7	33.4	43.8	42.0	32.5											
Genou + ss	37.5	45.6*	50.8*	49.3*	39.8	42.0	48.8*	47.7*	48.6*	50.3*	49.3*	49.3*	49.3*	44.0	34.1	44.0	34.1	2.8**	2.9**	9.9*	40.9	8.2*	8.4*											
Jagalene (P)+	37.0	41.9	48.1**	49.3*	37.0	40.4	44.7	43.3	45.1	46.5	45.8	45.4	45.4	44.0	34.1	44.0	34.1	2.8**	2.9**	9.9*	40.9	8.2*	8.4*											
Jerry	39.8	42.0	48.8*	49.3*	37.0	40.4	44.7	43.3	45.1	46.5	45.8	45.4	45.4	44.0	34.1	44.0	34.1	2.8**	2.9**	9.9*	40.9	8.2*	8.4*											
Judee ++ ss	37.5	45.6*	50.8*	49.3*	39.8	42.0	48.8*	47.7*	48.6*	50.3*	49.3*	49.3*	49.3*	44.0	34.1	44.0	34.1	2.8**	2.9**	9.9*	40.9	8.2*	8.4*											
Ledger (P)+	37.0	41.9	48.1**	49.3*	37.0	40.4	44.7	43.3	45.1	46.5	45.8	45.4	45.4	44.0	34.1	44.0	34.1	2.8**	2.9**	9.9*	40.9	8.2*	8.4*											
MT08172 (Colter)++	42.4	47.2*	50.5*	50.3*	42.0	38.5	44.5	43.4	45.8	47.1	47.3	47.3	47.3	42.8	40.7	45.7	47.9	47.1	45.3	42.4	43.4	45.3	45.3											
MT0871	40.7	44.2*	50.5*	50.3*	42.0	38.5	44.5	43.4	45.8	47.1	47.3	47.3	47.3	42.8	40.7	45.7	47.9	47.1	45.3	42.4	43.4	45.3	45.3											
MTCL1067	38.9	46.9*	51.4*	50.5*	42.0	38.3	43.3	42.6	43.5	45.2	44.8	45.1	45.1	32.0	23.3	27.4	25.8	25.0	26.8	26.7	26.7	26.7	26.7											
MTS0808 (Warhorse)++ ss	31.0	38.3	43.3	42.6	31.0	38.3	43.3	42.6	43.5	45.2	44.8	45.1	45.1	9.8*	8.1*	8.1**	8.2**	9.8**	9.2**	8.4**	8.3**	8.3**	8.3**											
MTS0819 ss	44.4	43.5*	51.7**	52.2**	44.4	43.5	43.3	42.6	43.5	45.2	44.8	45.1	45.1	40.9	8.2*	8.4*	8.4*	8.4*	8.4*	8.4*	8.4*	8.4*	8.4*											
Norris (P, CL)+	32.8	45.5*	52.5**	52.3**	32.8	45.5*	52.5**	50.3**	51.7**	52.3**	52.2**	51.7**	51.7**	42.6	33.9	39.7	42.6	41.0	44.1	44.6	46.0	46.0	46.0											
Overland +	37.1	42.9	48.1	48.5	37.1	42.9	48.1	46.8	48.0	49.0	48.5	48.2	48.2	31.2	24.8	30.7	33.0	33.6	34.2	33.1	33.1	33.1	33.1											
Pryor (P)+	ns	5.8	4.7	3.1	ns	5.8	4.7	4.0	3.7	3.4	3.1	2.8	2.8	27.8	16.9	15.3	11.6	10.7	10.1	9.4	9.1	9.1	9.1											
Rampart ss	13.0	10.7	9.2	8.9	13.0	10.7	9.2	9.2	9.1	8.9	8.9	8.7	8.7	54	56	47	38	38	38	39	41	41	41											
SY Clearstone 2CL (P)++																																		
WB-Quake (P) ++ ss																																		
Yellowstone +																																		
Average	37.1	42.9	48.1	46.8	37.1	42.9	48.1	46.8	48.0	49.0	48.5	48.2	48.2	31.2	24.8	30.7	33.0	33.6	34.2	33.1	33.1	33.1	33.1											
LSD (0.05)	ns	5.8	4.7	3.1	ns	5.8	4.7	4.0	3.7	3.4	3.1	2.8	2.8	27.8	16.9	15.3	11.6	10.7	10.1	9.4	9.1	9.1	9.1											
C.V. (%)	13.0	10.7	9.2	8.9	13.0	10.7	9.2	9.2	9.1	8.9	8.9	8.7	8.7	54	56	47	38	38	38	39	41	41	41											

\*\* = indicates highest value within a column  
\* = indicates varieties with values equal to highest variety within a column based on Fisher's protected LSD (p=0.05)  
(P) = Private Variety; + = Protected Variety; ++ = PVP Pending  
1/ = Dwarf Smut Resistant; (HWW) = Hard White Winter Wheat  
ss = solid-stemmed sawfly resistant variety

Table 13. Precipitation (top, in inches) and Average Monthly Temperature (bottom, °F) for Crop Year 2011-2012

Agricultural Research Center	Sept. 2011	Oct. 2011	Nov. 2011	Dec. 2011	Jan. 2012	Feb. 2012	Mar. 2012	Apr. 2012	May 2012	June 2012	July 2012	Aug 2012	Total Average
	Western Triangle, Conrad	0.30	1.84	0.18	0.12	0.17	0.02	0.53	1.04	1.49	2.46	0.82	1.32
	60.0	44.7	30.9	29.2	24.3	25.9	36.2	44.5	50.0	58.4	69.3	67.1	45.1
Northern, Havre	0.39	0.04	0.31	0.07	0.18	0.14	0.60	2.18	2.98	1.43	0.74	0.40	9.46
	59.9	47.1	30.9	29.4	22.4	25.3	39.0	46.6	52.0	62.3	72.7	68.7	46.4
Northwestern, Kalispell	0.91	2.46	0.46	0.40	1.08	1.15	1.16	1.35	2.11	7.11	1.41	0.56	20.16
	56.2	43.3	31.6	28.0	26.4	28.2	36.7	45.2	48.8	54.9	65.2	63.1	44.0
Central, Moccasin	0.46	0.88	0.57	0.17	0.60	0.22	0.85	2.64	1.56	1.47	0.83	0.74	10.99
	59.5	48.1	31.5	28.8	26.7	27.1	40.2	4545.4	49.2	59.1	71.3	67.5	46.2
Southern, Huntley	0.33	1.79	0.50	0.31	0.57	0.16	1.04	0.91	1.97	0.24	0.36	0.22	8.40
	62.9	51.3	34.1	31.9	27.4	29.7	44.5	49.7	54.6	66.8	76.6	70.8	50.1
Northeastern, Sidney	0.95	0.41	0.35	0.10	0.09	0.21	0.05	1.72	2.68	1.11	1.51	0.76	9.94
	59.6	49.7	31.3	26.4	24.1	22.5	43.6	48.5	55.7	67.0	75.5	68.4	47.7
Williston, N. Dakota	1.46	0.72	0.17	0.07	0.80	0.17	0.11	1.65	2.12	2.39	2.41	0.77	12.84
	61.3	49.7	30.5	25.3	22.2	22.7	43.1	47.7	54.9	65.9	76.2	70.0	47.4
Post Farm, Bozeman	0.66	0.03	0.64	0.17	0.17	0.44	1.07	2.48	1.77	0.72	0.58	0.27	9.00
	61.6	47.8	31.7	26.3	30.1	27.9	41.6	47.3	50.7	60.6	70.5	68.6	47.1

**Table 14. Selected agronomic characters, cereal quality evaluations and disease reactions of hard winter wheat varieties.**

Variety	Agronomic Characters						Cereal Quality			Disease Reactions <sup>8/</sup>			
	Maturity <sup>1/</sup>	Chaff Color	Winter Survival <sup>2/</sup>	Straw Strength <sup>3/</sup>	Stem solid <sup>4/</sup>	Coleoptile length <sup>5/</sup>	Milling <sup>6/</sup>	Baking <sup>6/</sup>	PPO <sup>7/</sup>	Dwarf Smut	Stripe Rust	Stem Rust	Leaf Rust
Accipiter	M-L	White	5	MS		M	2	3	H	S	S	MR	R
AP 503 CL2	M	White	2	S		M	3	5	H	S	R	MR	S
Art	E	White	2	S		S	3	2	M	S	MR	MR	M
Bearpaw	M	White	2	M	21	M	4	3	H	S	S	R	S
Broadview	M	White	5	S		S	3	3	H	S	S	R	R
Bynum	M	Brown	2	M	18	L	5	4	M	S	R	MS	S
Carter	M	White	3	S	14	S	4	5	M	S	S	MS	M
CDC Falcon	M	White	4	S	7	S	3	3	H	S	S	MR	R
Colter (MT08172)	M	White	3	S		S	3	4	M	S	R	R	S
Curlew	M	Brown	2	S		M	4	3	L	R	R	VS	S
Decade	M	White	4	S		M	3	4	H	S	S	R	MS
Genou	M	White	2	M	18	M	4	4	H	S	S	S	MR
Jagalene	E	White	2	S		M	4	3	H	S	R	MR	MS
Jerry	M	White	5	M		M	3	3	H	S	S	R	R
Judee	M	White	2	S	19	M	3	4	H	S	R	S	S
Ledger	M	White	2	S	10	M	5	3	M-H	S	S	S	MS
McGill	E	White	3	S		M	2	3	M	S	S	R	-
Norris	E	White	3	S		M	3	3	M	S	S	S	MS
Overland	E	White	3	S		M	3	2	M	S	S	R	R
Peregrine	M-L	White	5	MS		M	3	3	M	S	R	MR	R
Promontory	M	Brown	2	MS		S	4	3	L	R	R	VS	S
Pryor	M	White	3	S		S	3	2	H	S	S	S	S
Radiant	M-L	White	4	S		S	3	3	H	S	R	VS	M
Rampart	M	Brown	2	MW	21	L	4	5	M	S	R	MR	S
Robidoux	E	White	2	S		S	3	3	H	S	MS	S	S
SY Clearstone 2CL	M	White	3	S		S	3	3	M	S	R	MR	-
SY Wolf	M	White	3	S		M	3	2	M	S	R	R	-
WB-Quake	M-L	White	3	S	19	M	4	4	H	S	R	MR	MR
Warhorse (MTS080)	M	White	3	S	21	M	3	3	H	S	R	R	MR
Yellowstone	M	White	4	S		S	3	4	M	S	R	S	MS

1/ VE = Very Early, E = Early, M = Medium, L = Late, VL = Very Late

2/ 5 = Best Winter survival (over several years at Sidney, Williston and Moccasin)

3/ W = Weak

5/ L = long

6/ 5 = Superior

7/ PPO = Polyphenol Oxidase

MW = Medium Weak

M = medium

4

(low is better for noodles)

M = Medium

S = short

3

L = low

MS = Medium Strong

- = no info.

2

M = medium

S = Strong

1 = Inferior

H = high

4/ scored 5-25, 25 = most solid

Combined Bozeman, Conrad, Havre, Moccasin, and Sidney data; 2010-2012

varieties with no number were not evaluated

8/ R = Resistant

MR = Moderately Resistant

M = Moderate

MS = Moderately Susceptible

S = Susceptible

VS = Very Susceptible

- = no information

## Additional Descriptive Information for Winter Wheat Varieties

### New for the 2013 Bulletin:

**Colter** (MT08172) – is an awned, white glumed, high yielding hard red winter wheat to be released in fall 2013 by the Montana Agricultural Experiment Station. Colter is similar to Yellowstone for grain yield and most agronomic traits with the exception that Colter is about 0.5 lb/bu higher for test weight and has superior stem rust resistance relative to Yellowstone. Colter is moderately resistant to stripe rust, but susceptible to leaf rust. Colter has excellent milling and baking bread quality, similar to Yellowstone. PVP, Title V will be applied for.

**SY Clearstone 2CL** (MTCL1077) – a 2-gene CLEARFIELD hard red winter wheat developed by Montana Agricultural Experiment Station in 2012 and licensed exclusively to Syngenta Seeds. SY Clearstone wheat 2CL is very similar to Yellowstone. It is a medium maturing, medium tall, white chaffed wheat with average winter hardiness. It is a high yielding wheat with average test weight and protein. SY Clearstone 2CL is resistant to stripe rust and has moderate resistance to stem rust, the latter an improvement over Yellowstone. SY Clearstone 2CL is a medium PPO variety with average mill and above average bake properties. PVP, Title V will be applied for. Additionally, the CLEARFIELD genes are patented.

**Warhorse** (MTS0808) - is an awned, white glumed, solid-stemmed hard red winter wheat to be released in fall 2013 by the Montana Agricultural Experiment Station. Warhorse has medium maturity and has medium short, semi-dwarf height. Stem solidness is similar to that of Bearpaw and Rampart, while sawfly cutting of stems is very low (similar to Rampart). Warhorse yield is similar to Judee, while test weight and protein are above average. Warhorse is resistant to both stem and stripe rust. Warhorse has acceptable mill and bake qualities. PVP, Title V will be applied for.

### Varieties previously in bulletin:

**Accipiter** – hard red winter wheat developed by the Crop Development Center, Saskatoon, Saskatchewan and registered in 2008. Accipiter is a medium to late maturing average height wheat with white chaff. Accipiter has above average yield, below average test weight and protein, and excellent winter hardiness. Accipiter is susceptible to stripe rust and moderately resistant to stem rust. Accipiter has below average milling and average

baking quality. PVP, Title V is pending (Certificate #201100370).

**AP503 CL2** – a 2-gene CLEARFIELD hard red winter wheat released by AgriPro in 2007. AP503 CL2 is a medium maturing short wheat with white chaff. AP503 CL2 has below average yield, above average test weight, average protein, and below average winter hardiness. AP503 CL2 appears resistant to stripe rust and moderately resistant to stem rust. AP503 CL2 has average milling and above average baking quality. PVP, Title V has been issued (Certificate #200800322). Additionally, the CLEARFIELD genes are patented.

**Art** – hard red winter wheat developed by Syngenta (AgriPro) Seeds in 2007. Art is an early maturing, short statured wheat, with white chaff. Art has average yield, above average test weight and protein, and below average winter hardiness. Art is moderately resistant to stripe and stem rust. Art has average milling and below average baking quality. PVP, Title V has been issued (Certificate #200700349).

**Bearpaw** – hard red winter wheat developed by the Montana Agricultural Experiment Station in 2011. Bearpaw is a white-glumed, solid-stem, semi-dwarf (*Rht1*) wheat with medium maturity. Bearpaw has average yield, test weight, and protein, and below average winter hardiness. Bearpaw is resistant to prevalent races of stem rust but susceptible to stripe and leaf rust. Stem-solidness of Bearpaw is most similar to Rampart. Bearpaw is a high PPO variety with above average milling and average baking properties. PVP, Title V is pending (Certificate #201200407).

**Broadview** – hard red winter wheat developed by the Lethbridge, Alberta winter wheat breeding program in 2009. Broadview is a medium maturing, medium statured wheat, with white chaff. Broadview has above average yield, average test weight and protein, and excellent winter hardiness. Broadview is susceptible to stripe rust and resistant to stem rust. Broadview is a high PPO variety with average milling and baking properties.

**Bynum** – a CLEARFIELD (CL) wheat with imidazolinone tolerance, developed by the Montana Agricultural Experiment Station in 2005 and licensed to WestBred LLC. Bynum is a solid stem “Rampart-type” CLEARFIELD hard red winter cultivar similar in most characteristics to Rampart.

It is lower yielding than Norris and similar in yield to MT1159CL. Bynum has a solid stem, high grain protein, and excellent bread baking quality. Bynum is resistant to stripe rust and has some resistance to stem rust. PVP, Title V has been issued (Certificate #200600285). Additionally, the CLEARFIELD gene is patented.

**Carter** – a semi-solid stem hard red winter wheat released by WestBred LLC in 2007. Carter is a medium maturity semidwarf wheat. It has average yield, test weight, and winterhardiness and good protein. Carter is moderately susceptible to stem rust and susceptible to stripe rust. Carter has above average milling and baking quality. PVP, Title V has been issued (Certificate #200800383).

**CDC Falcon** – hard red winter wheat developed by the Crop Development Center, Saskatoon, Saskatchewan and registered in 1998. Licensed to WestBred LLC. Superior stem and leaf rust resistance over all current winter wheat varieties in western Canada. High yield, good winterhardiness, semidwarf, short strong straw, especially good for direct seeding and straight cut harvest. CDC Falcon is moderately resistant to stem rust and susceptible to stripe rust. It is rated as having acceptable milling and baking quality. CDC Falcon is protected under the Plant Variety Protection Act, but not the Title V option (Certificate #200800322).

**Curlew** – hard red winter wheat released by Utah in 2009. Curlew is an early to medium maturing tall wheat with brown chaff. In the initial year of testing in Montana, Curlew had above average yield and test weight and average protein. Curlew appears resistant to stripe rust but very susceptible to stem rust. Curlew is resistant to dwarf bunt. Curlew has above average milling and average baking properties. It is a low PPO variety.

**Decade** – hard red winter wheat developed by the Montana Agricultural Experiment Station and released jointly with North Dakota (pending at publication) in 2010. Decade is an early to medium maturing reduced height wheat with white chaff. Decade is a high yielding wheat with good winterhardiness and medium to high test weight and protein. Decade is resistant to prevalent races of stem and stripe rust. Decade has excellent milling and baking quality. Seed available fall 2010. PVP, Title V has been issued (Certificate #201100096).

**Genou** – a solid-stem hard red winter wheat with improved yield potential and cold tolerance relative to Rampart. Stem solidness is relatively good, although not as good as Rampart. Test weight, maturity, plant height, grain protein, and end-use

qualities are similar to those of Rampart and Vanguard. Genou is susceptible to both stem and stripe rust. Foundation seed was made available in fall of 2004. Genou (French for knee) is named after a school house in The Knees area of Chouteau County. PVP with Title V option has been issued (Certificate #200500334).

**Jagalene** – Developed from the cross “Abilene/Jagger” and released by AgriPro in 2003. Jagalene is a hollow-stemmed hard red winter variety. It has been tested in Montana Intrastate Trials and Off-Station trials during the 2003 through 2005 seasons. It has exhibited consistently high yields in Crop Reporting Districts 1 through 5. Jagalene has excellent test weight ranking higher than all checks. It is a semi-dwarf with height shorter than all checks except CDC Falcon. Lodging resistance is very good. It has early heading, nearly 6 days earlier than Neeley and earlier than all checks. Winterhardiness levels would be considered average to below average. It has excellent general disease resistance, including the entire soil virus complex, stem rust, stripe rust, tan spot and septoria. Protein is average compared to the checks. Jagalene has excellent milling and very good baking characteristics. This variety is protected under the Plant Variety (Certificate #200200160) Protection Act and can only be sold or advertised by variety name as a class of certified seed.

**Jerry** – hard red winter wheat released by North Dakota State University in 2001. It is white-chaffed and awned and similar in maturity to Roughrider. Jerry has good winter hardiness and is a top yielder in areas where winterkill can occur. Jerry has average test weight and protein under Montana conditions. It has good resistance to prevalent races of stem and leaf rust, but is susceptible to stripe rust. Mixing properties and baking performance are equal to Roughrider.

**Judee** – hard red winter wheat developed by the Montana Agricultural Experiment Station in 2011. Judee is a white-glumed, solid-stem, semi-dwarf (*Rht1*) wheat with medium maturity. Judee has average yield, test weight, and protein, and below average winter hardiness. Judee is susceptible to prevalent races of stem and leaf rust but resistant to stripe rust. Stem-solidness of Judee is most similar to Genou. Judee is a high PPO variety with average mill and above average bake properties. PVP, Title V is pending (Certificate #201200161).

**Ledger** – hard red winter wheat developed by WestBred LLC and released in 2004. Ledger is an early maturing wheat, semidwarf wheat with



average winter hardiness. The Montana Intrastate Winter Wheat Program testing shows this variety to be of average yield and protein with above average test weight. Ledger is susceptible to stripe rust and stem rust. Milling and baking characteristics are acceptable. Ledger is protected under the Plant Variety Protection Act, but not the Title V option (Certificate #200600063).

**McGill** – hard red winter wheat developed by Nebraska in 2010. McGill is an early maturing, medium statured wheat with white glumes. McGill has above average yield, average test weight, and below average protein. Winter-hardiness was above average in 2011 at Sidney. McGill is susceptible to stripe rust, but resistant to stem rust. McGill has below average milling and average baking properties. PVP, Title V has been issued (Certificate #201100399).

**Norris** - a CLEARFIELD (CL) wheat with imidazolinone tolerance, developed by the Montana Agricultural Experiment Station in 2005 and licensed to WestBred LLC. Norris is a high yielding hard red winter CLEARFIELD cultivar that could replace MT1159CL once seed becomes available. Norris is significantly higher in yield than other CLEARFIELD checks and similar in yield to Neeley. Norris has high test weight, good crop tolerance to herbicide, and is relatively early in heading compared to Montana varieties. Grain protein of Norris is relatively low, but milling and baking characteristics are acceptable. Norris is susceptible to both stem and stripe rust. PVP, Title V has been issued (Certificate #200600286). Additionally, the CLEARFIELD gene is patented.

**Overland** – hard red winter wheat developed in Nebraska and released jointly with South Dakota in 2007. Overland is an early maturing average height wheat with white chaff. In the initial year of testing in Montana, Overland had average yield, test weight, and protein. Overland is resistant to stem rust and susceptible to stripe rust. Overland has average milling and below average baking quality. PVP, Title V has been issued (Certificate #200700333).

**Peregrine** – hard red winter wheat developed by the Crop Development Center, Saskatoon, Saskatchewan and registered in 2008. Peregrine is a medium to late maturing tall wheat with white chaff. Peregrine has average yield, above average test weight, below average protein, and good winter hardiness. Peregrine appears resistant to stripe rust and moderately resistant to stem rust. Peregrine has average milling and baking quality. PVP, Title V is pending (Certificate #201100371).

**Promontory** – Released by the Utah Agricultural Experiment Station in 1991. It is a hard red winter wheat of medium height with awns and bronze chaff. Promontory is a high yielding line with excellent test weight. It has poor winterhardiness. Promontory is resistant to dwarf bunt and stripe rust and susceptible to stem rust. Promontory has average milling and above average baking characteristics. It has low PPO and could be used as a dual-purpose (bread and noodles) variety.

**Pryor** – hard red winter wheat released by Western Plant Breeders in 2002. Pryor is a white chaffed, awned variety with short stature and medium winter hardiness. Pryor is a high yielding variety with average test weight and below average protein. It is susceptible to stem rust and stripe rust and moderately resistant to leaf spot complex. Pryor has average milling and below average baking characteristics. This variety is protected under the Plant Variety Protection Act without the Title V option (Certificate #200400072).

**Radiant** – hard red winter wheat released by Alberta in 2002 and marketed by Meridian Seeds. Radiant is a medium to late maturing, medium tall wheat with white chaff. Radiant has good winter hardiness in North Dakota tests. In the initial year of testing in Montana, Radiant had average yield, average test weight, and below average protein. Radiant appears resistant to stripe rust and very susceptible to stem rust. Radiant has average milling and baking quality.

**Rampart** – Released by the Montana Agricultural Experiment Station in 1996. It is an awned, red chaffed, solid-stemmed hard red winter wheat variety. The kernel is long with a sloping back and a heavy brush. The cheeks are rounded to angular with an open crease. Rampart is resistant to the wheat stem sawfly. It is moderately resistant to prevalent races of stem rust. Rampart is resistant to stripe rust. It is susceptible to leaf rust, dwarf smut and the Russian wheat aphid. Rampart has excellent milling and baking properties and is a sister line to Vanguard.

**Robidoux** – hard red winter wheat developed by Nebraska in 2010. Robidoux is an early maturing, medium statured wheat, with white chaff. Robidoux has above average yield, average test weight, and below average protein, and winter hardiness. Robidoux is moderately susceptible to stripe rust and susceptible to stem rust. Robidoux is a high PPO variety with average mill and bake qualities. PVP, Title V is pending (Certificate #201100398).

**SY-Wolf** – hard red winter wheat developed by Syngenta (AgriPro) Seeds in 2010. SY-Wolf is a medium maturing, short statured wheat with white glumes. SY-Wolf has above average yield and test weight and average protein. Winter-hardiness was average in 2011 at Sidney. SY-Wolf is moderately susceptible to moderately resistant (MS/MR) to stripe rust, but resistant to stem rust. Boomer has average milling and below average baking properties. PVP, Title V has been issued (Certificate #201100390).

**WB-Quake** – hard red winter wheat developed by WestBred (Monsanto) in 2011. WB-Quake is a medium to late maturing, medium statured solid-stemmed wheat, with white chaff. WB-Quake has above average yield, average test weight and protein with average winter hardiness. WB-Quake is resistant to stripe rust and moderately resistant to stem rust. WB-Quake is a high PPO variety with above average milling and baking properties. PVP, Title V is pending (Certificate #201100471).

**Yellowstone** – hard red winter wheat developed by the Montana Agricultural Experiment Station and released to seed growers in 2005. Yellowstone is a very high yielding winter hardy variety with medium test weight, maturity, height, and grain protein. Yellowstone has excellent baking and good Asian noodle quality. It is moderately resistant to TCK smut and resistant to stripe rust, but susceptible to stem rust. Yellowstone potentially could occupy acreage currently planted to Neeley, Tiber CDC Falcon, Paul, Promontory, and Morgan. PVP, Title V has been issued (Certificate #200600284).

## Plant Variety Protection

The Plant Variety Act, signed into law in 1970, offers legal protection to developers of new varieties of plants which reproduce sexually – that is, through seeds. The law provides for a Plant Variety Protection Office in the U.S. Department of Agriculture. The office receives and processes applications and when “novelty” is established, issues a certificate granting protection rights specified by the applicant.

The owner (or developer) holding a “certificate of protection” has complete control over the variety for 20 years. The law provides two types of protection:

### 1. Without Seed Certification

The owner of the protected variety may exclude others from reproducing the variety, selling it, offering it for sale, importing or exporting it, or use it

in the commercial production of a hybrid or a different variety without permission. In this sense, the owner of a protected variety may bring civil damage action against anyone who infringes upon his rights.

### 2. Certified Seed Option

The owner may specify that the seed of his variety “...be sold or advertised only as a class of Certified Seed”. Production and sale of such seed by variety name, when not certified, constitute a violation of the Federal Seed Act. This means of protection may be used extensively for publicly as well as privately developed varieties.

Amendments to the Plant Variety Protection Act (PVPA) have passed both houses of Congress and been signed into law by the President. These amendments went into effect in 1995. The farmers exemption has been changed for new varieties. Seed for varieties issued a certificate after April 4, 1995, may only be purchased from the owner or his agent. A farmer can only save seed of these varieties for use on his own farm and cannot sell seed of the protected variety to his neighbor.

A variety protected under the certification option does not permit a farmer producing seed to sell or offer for sale or advertise by variety name unless it is certified. Sale of such seed by variety name as uncertified seed will constitute a violation of the Federal Seed Act. Interstate movement of seed is subject to inspection by Federal Seed Control officials. Seed within the state is subject to inspection by State Department of Agriculture inspectors.

Owners of protected varieties will give public notice that their variety is protected by affixing to the label or container the words: “Unauthorized Propagation Prohibited” or the words, “Unauthorized Seed Multiplication Prohibited”. Producers must check the label (tag) or the container for the above wording

Publication reviewed and/or data supplied by the following Montana and North Dakota research staff:

Mr. Jim Berg, Research Associate, Plant Sciences and Plant Pathology Department, Montana State University, Bozeman, Montana.

Dr. Phil Bruckner, Professor, Winter Wheat Breeding, Plant Sciences and Plant Pathology Department, Montana State University, Bozeman, Montana.

Dr. Gerald W. Bergman, Superintendent, Williston Research and Extension Center, North Dakota State University, Williston, ND

Mr. Benri Deanon, Research Associate, Central Agricultural Research Center, Moccasin, Montana.

Dr. Alan Dyer, Assistant Professor, Plant Sciences and Plant Pathology Department, Montana State University, Bozeman, Montana.

Dr. Joyce Eckhoff, Associate Professor of Agronomy, Eastern Agricultural Research Center, Sidney, Montana.

Dr. Bill Grey, Adjunct Assistant Professor and Montana Foundation Seed Stocks Manager, Plant Sciences and Plant Pathology Department, Montana State University, Bozeman, Montana.

Dr. Ken Kephart, Superintendent and Associate Professor of Agronomy, Southern Agricultural Research Center, Huntley, Montana

Ms. Peggy Lamb, Agronomist, Northern Agricultural Research Center, Havre, Montana.

Mr. Ron Larson, Manager, Montana Seed Growers Association, Montana State University, Bozeman, Montana.

Ms. Sara Loomer, Research Specialist, Williston Research and Extension Center, North Dakota State University, Williston, ND

Mr. John Miller, Research Associate, Western Triangle Research Center, Conrad, Montana.

Ms. Deanna Nash, Cereal Quality Laboratory Manager, Plant Sciences and Plant Pathology Department, Montana State University, Bozeman, Montana.

Mr. Mat Peterson-Walter, Research Specialist, Southern Agricultural Research Center, Huntley, Montana

Dr. Gaddi V. P. Reddy, Superintendent, Western Triangle Research Center, Conrad, Montana.

Dr. Robert Stougaard, Superintendent and Professor of Weed Science, Northwestern Agricultural Research Center, Kalispell, Montana.

Mr. Dave Wichman, Superintendent and Associate Professor of Agronomy, Central Agricultural Research Center, Moccasin, Montana.

**Note: Information in this article is available on the web at:**

**<http://plantsciences.montana.edu/crops>**